



Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin no.-62 for the Rajkot district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 2nd To 6th November- 2013)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	0.0
Maximum temperature (°C)	33.5-34.0
Minimum temperature (°C)	17.0-18.5
Relative Humidity (%) I	69-94
Relative Humidity (%) II	29-34
Wind speed (Kmph)	1.58-2.32

Weather forecast until 08-30 hrs of 06/11/2013

(1) RAJKOT DISTRICT:

	02/ 11/13	03/ 11/13	04/ 11/13	05/ 11/13	06/ 11/13
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	33	34	33	34	34
Min.Temp.(°C)	20	20	21	21	21
Total CC (octa)	1	0	0	0	1
RH-I (%)	58	51	40	36	45
RH-II (%)	32	29	26	21	20
Wind Speed (kmph)	5	7	10	6	9
Wind Direction(deg.)	157	87	38	26	342

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none"> ➤ Incoming five days sky will remain mostly clear. Maximum temperature will raise up to 34 °c in day time. ➤ Farmers are also advised that it is proper time for sowing of garlic, gram and mustard. ➤ Apply irrigation and nitrogenous fertilizer to cotton and vegetables.
-----------------------	--

Crops	Stage	Insect/Pest/Variety	Advisory
Garlic	Sowing 	Gujarat Garlic-4 or Local	Carry out sowing of Garlic, If kharif crop is harvested.
Chick pea	Sowing	Gujarat chana-2 (Unirrigated), Gujarat chana-3 (irrigated)	Carry out sowing of Gram, If kharif crop is harvested.
Mustard	Sowing	Varuna, Gujarat rai-3	Carry out sowing of Mustard, If kharif crop is harvested.
Cotton	Squaring, flowering & ball formation & opening	Jassid/Thrips/Aphid →   Angular leaf spot, Alternaria leaf spot  	Acetamiprid 20 % SP@ 7 g /10 lit of water is Two Spray of Streptocyclin 1 g + Copper oxychloride 40 g/10 lit of water at 10 days interval starting from the initiation of disease.

		Fertilizer & Irrigation →	Spray NPK (19-19-19) and micronutrient 50 g/10 lit of water for quick regrowth.
Castor	Vegetative & reproductive	Jassid, Whitefly →	Spray Thiomethoxam 25 % WG @ 4 g / 10 lit of water
Onion	Growing	Purple blotch →  Thrips 	Alternative spray of Mencozeb 25 gm and hexaconazole 15 ml in 10 liters of water at 10 days interval. Spray Profenophos 20 ml/ 10 liters of water.
Vegetables	Growing/fruiting	Sucking pest Fruit borer  	Current weather is favourable for outbreak of sucking pest and disease. To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water and hexaconazole 5% SC @ 10 ml/10 lit of water. In brinjal, cauliflower, cabbage, tomato etc spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer, diamond back moth in cloud free days. Spray NPK (19-19-19) and micronutrient 50 g/10 lit of water for quick regrowth.
Chili	Flowering & fruiting	Thrips, →  Wilting →	Alternative spray of 20 ml fipronil, 20 ml profenophos and 10 ml of DDVP at 10 days interval. Drenching of Trichoderma viridi 4 gm in 1 liter of water.
Lady's finger	Flowering/ fruiting	Shoot and fruit borer → Mites →	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 liters of water at 15 days interval(4 spray). Spraying of Dicofol @ 2 ml/liters of water is advised.
Runner vegetable	Flowering/ fruiting	Downy mildew	Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval.
Custard apple	Fruit development	Mealy bug- 	Spray quinalphos 20 ml/10 litres of water.
Livestock & poultry 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick and for prevention of foot & mouth disease in animals do vaccination with FMD vaccine. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid flys and mosquitoes.		



**Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523**



Agro met Advisory Service bulletin for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 2nd To 6th November - 2013)

Weather forecast until 08-30 hrs of 06/11/2013

(2) JAMNAGAR DISTRICT:

	02/ 11/13	03/ 11/13	04/ 11/13	05/ 11/13	06/ 11/13
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	30	31	30	31	31
Min.Temp.(°C)	21	22	22	22	22
Total CC (octa)	1	0	0	0	0
RH-I (%)	68	52	39	34	56
RH-II (%)	43	39	30	24	24
Wind Speed (kmph)	7	10	12	8	11
Wind Direction(deg.)	225	166	47	81	336

Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none"> ➤ Incoming five days sky will remain mostly clear. Maximum temperature will raise up to 31 °c in day time. ➤ Farmers are also advised that it is proper time for sowing of garlic, gram and mustard. ➤ Apply irrigation and nitrogenous fertilizer to cotton and vegetables. 		
Crops	Stage	Insect/Pest/Variety	Advisory
Garlic	Sowing	Gujarat Garlic-4 or Local	Carry out sowing of Garlic, If kharif crop is harvested.
Chick pea	Sowing	Gujarat chana-2 (Unirrigated), Gujarat chana-3 (irrigated)	Carry out sowing of Gram, If kharif crop is harvested.
Mustard	Sowing	Varuna, Gujarat rai-3	Carry out sowing of Mustard, If kharif crop is harvested.
Cotton	Squaring, flowering & ball formation & opening	Jassid/Thrips/Aphid →   Angular leaf spot, Alternaria leaf spot   Fertilizer & Irrigation →	Acetamiprid 20 % SP@ 7 g /10 lit of water is Two Spray of Streptocyclin 1 g + Copper oxychloride 40 g/10 lit of water at 10 days interval starting from the initiation of disease. Spray NPK (19-19-19) and micronutrient 50 g/10 lit of water for quick regrowth.
Castor	Vegetative & reproductive	Jassid, Whitefly →	Spray Thiomethoxam 25 % WG @ 4 g / 10 lit of water
Onion	Growing	Purple blotch →	Alternative spray of Mencozeb 25 gm and

		 Thrips 	hexaconazole 15 ml in 10 liters of water at 10 days interval. Spray Profenophos 20 ml/ 10 liters of water.
Vegetables	Growing/fruiting	Sucking pest Fruit borer  	Current weather is favourable for outbreak of sucking pest and disease. To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water and hexaconazole 5% SC @ 10 ml/10 lit of water. In brinjal, cauliflower, cabbage, tomato etc spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer, diamond back moth in cloud free days. Spray NPK (19-19-19) and micronutrient 50 g/10 lit of water for quick regrowth.
Chili	Flowering & fruiting	Thrips, →  Wilting →	Alternative spray of 20 ml fipronil, 20 ml profenophos and 10 ml of DDVP at 10 days interval. Drenching of Trichoderma viridi 4 gm in 1 liter of water.
Lady's finger	Flowering/ fruiting	Shoot and fruit borer → Mites →	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 liters of water at 15 days interval(4 spray). Spraying of Dicofol @ 2 ml/liters of water is advised.
Runner vegetable	Flowering/ fruiting	Downy mildew	Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval.
Custard apple	Fruit development	Mealy bug- 	Spray quinalphos 20 ml/10 litres of water.
Livestock & poultry 		Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick and for prevention of foot & mouth disease in animals do vaccination with FMD vaccine. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid flys and mosquitoes.	

IAAS Unit, DFRS, JAU, Targhadia



**Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523**



Agro met Advisory Service bulletin for the Amreli district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 2nd October To 6th November - 2013)

Weather forecast until 08-30 hrs of 06/11/2013

(3) AMRELI DISTRICT:

	02/ 11/13	03/ 11/13	04/ 11/13	05/ 11/13	06/ 11/13
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	32	33	32	33	33
Min.Temp.(°C)	19	20	20	20	20
Total CC (octa)	1	0	0	0	1
RH-I (%)	55	53	44	40	46
RH-II (%)	33	32	31	27	27
Wind Speed (kmph)	5	8	11	7	7
Wind Direction(deg.)	78	43	32	34	300

Agro meteorological Advisories

General Advice	<ul style="list-style-type: none"> ➤ Incoming five days sky will remain mostly clear. Maximum temperature will raise up to 33 °c in day time. ➤ Farmers are also advised that it is proper time for sowing of garlic, gram and mustard. ➤ Apply irrigation and nitrogenous fertilizer to cotton and vegetables.
-----------------------	--

Crops	Stage	Insect/Pest/Variety	Advisory
Garlic	Sowing	Gujarat Garlic-4 or Local	Carry out sowing of Garlic, If kharif crop is harvested.
Chick pea	Sowing	Gujarat chana-2 (Unirrigated), Gujarat chana-3 (irrigated)	Carry out sowing of Gram, If kharif crop is harvested.
Mustard	Sowing	Varuna, Gujarat rai-3	Carry out sowing of Mustard, If kharif crop is harvested.
Cotton	Squaring, flowering & ball formation & opening	Jassid/Thrips/Aphid →   Angular leaf spot, Alternaria leaf spot   Fertilizer & Irrigation →	Acetamiprid 20 % SP@ 7 g /10 lit of water is Two Spray of Streptocyclin 1 g + Copper oxychloride 40 g/10 lit of water at 10 days interval starting from the initiation of disease. Spray NPK (19-19-19) and micronutrient 50 g/10 lit of water for quick regrowth.
Castor	Vegetative & reproductive	Jassid, Whitefly →	Spray Thiomethoxam 25 % WG @ 4 g / 10 lit of water
Onion	Growing	Purple blotch →	Alternative spray of Mencozeb 25 gm and

		 Thrips 	hexaconazole 15 ml in 10 liters of water at 10 days interval. Spray Profenophos 20 ml/ 10 liters of water.
Vegetables	Growing/fruiting	Sucking pest Fruit borer  	Current weather is favourable for outbreak of sucking pest and disease. To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water and hexaconazole 5% SC @ 10 ml/10 lit of water. In brinjal, cauliflower, cabbage, tomato etc spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer, diamond back moth in cloud free days. Spray NPK (19-19-19) and micronutrient 50 g/10 lit of water for quick regrowth.
Chili	Flowering & fruiting	Thrips, →  Wilting →	Alternative spray of 20 ml fipronil, 20 ml profenophos and 10 ml of DDVP at 10 days interval. Drenching of Trichoderma viridi 4 gm in 1 liter of water.
Lady's finger	Flowering/ fruiting	Shoot and fruit borer → Mites →	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 liters of water at 15 days interval(4 spray). Spraying of Dicofol @ 2 ml/liters of water is advised.
Runner vegetable	Flowering/ fruiting	Downy mildew	Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval.
Custard apple	Fruit development	Mealy bug- 	Spray quinalphos 20 ml/10 litres of water.
Livestock & poultry 		Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick and for prevention of foot & mouth disease in animals do vaccination with FMD vaccine. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid flys and mosquitoes.	

IAAS Unit, DFRS, JAU, Targhadia



Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 2nd October To 6th November - 2013)

Weather forecast until 08-30 hrs of 06/11/2013

(4) SURENDRANAGAR DISTRICT:

	02/ 11/13	03/ 11/13	04/ 11/13	05/ 11/13	06/ 11/13
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	33	32	33	33	33
Min.Temp.(°C)	22	23	23	23	22
Total CC (octa)	0	0	0	0	0
RH-I (%)	53	46	32	29	37
RH-II (%)	29	29	21	16	16
Wind Speed (kmph)	4	9	11	5	7
Wind Direction(deg.)	162	171	32	100	325

Agro meteorological Advisories

General Advice	<ul style="list-style-type: none"> ➤ Incoming five days sky will remain mostly clear. Maximum temperature will raise up to 33 °c in day time. ➤ Farmers are also advised that it is proper time for sowing of garlic, gram and mustard. ➤ Apply irrigation and nitrogenous fertilizer to cotton and vegetables.
-----------------------	--

Crops	Stage	Insect/Pest/Variety	Advisory
Garlic	Sowing	Gujarat Garlic-4 or Local	Carry out sowing of Garlic, If kharif crop is harvested.
Chick pea	Sowing	Gujarat chana-2 (Unirrigated), Gujarat chana-3 (irrigated)	Carry out sowing of Gram, If kharif crop is harvested.
Mustard	Sowing	Varuna, Gujarat rai-3	Carry out sowing of Mustard, If kharif crop is harvested.
Cotton	Squaring, flowering & ball formation & opening	Jassid/Thrips/Aphid →   Angular leaf spot, Alternaria leaf spot   Fertilizer & Irrigation →	Acetamiprid 20 % SP@ 7 g /10 lit of water is Two Spray of Streptocyclin 1 g + Copper oxychloride 40 g/10 lit of water at 10 days interval starting from the initiation of disease. Spray NPK (19-19-19) and micronutrient 50 g/10 lit of water for quick regrowth.
Castor	Vegetative & reproductive	Jassid, Whitefly →	Spray Thiomethoxam 25 % WG @ 4 g / 10 lit of water
Onion	Growing	Purple blotch →	Alternative spray of Mencozeb 25 gm and

		 Thrips 	hexaconazole 15 ml in 10 liters of water at 10 days interval. Spray Profenophos 20 ml/ 10 liters of water.
Vegetables	Growing/fruiting	Sucking pest Fruit borer  	Current weather is favourable for outbreak of sucking pest and disease. To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water and hexaconazole 5% SC @ 10 ml/10 lit of water. In brinjal, cauliflower, cabbage, tomato etc spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer, diamond back moth in cloud free days. Spray NPK (19-19-19) and micronutrient 50 g/10 lit of water for quick regrowth.
Chili	Flowering & fruiting	Thrips, →  Wilting →	Alternative spray of 20 ml fipronil, 20 ml profenophos and 10 ml of DDVP at 10 days interval. Drenching of Trichoderma viridi 4 gm in 1 liter of water.
Lady's finger	Flowering/ fruiting	Shoot and fruit borer → Mites →	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 liters of water at 15 days interval(4 spray). Spraying of Dicofol @ 2 ml/liters of water is advised.
Runner vegetable	Flowering/ fruiting	Downy mildew	Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval.
Custard apple	Fruit development	Mealy bug- 	Spray quinalphos 20 ml/10 litres of water.
Livestock & poultry 		Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick and for prevention of foot & mouth disease in animals do vaccination with FMD vaccine. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid flys and mosquitoes.	

IAAS Unit, DFRS, JAU, Targhadia



હવામાન આગાહી આધારીત કૃષિ સલાહ બુલેટીન
ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ
તા. ૦૨-૧૧-૨૦૧૩ થી તા. ૦૬-૧૧-૨૦૧૩



(સંક્લીત કૃષિ મૈસમ સલાહ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડીયા અને
ભારત મૈસમ વિભાગ દ્વારા સંપન્ન)

કૃષિ હવામાન બુલેટીન નં : ૫૨/૨૦૧૩

પાછળા અઠવાડીયાનું હવામાન

હવામાન પરિબળો	ઉત્તર સૌરાષ્ટ્ર
	રાજકોટ
વરસાદ (મી.મી)	0.0
મહત્વમાન તાપમાન (સે.ગ્રે)	33.5-34.0
લઘૃતમ તાપમાન (સે.ગ્રે)	17.0-18.5
ભેજ (%) સવાર	69-94
બપોરબાદ	29-34
પવનની ગતિ (કિમી/કલાક)	1.58-2.32

હવામાન આગાહી:

૧. રાજકોટ જીલ્લો

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૦૨-૧૧-૧૩	૦૩-૧૧-૧૩	૦૪-૧૧-૧૩	૦૫-૧૧-૧૩	૦૬-૧૧-૧૩
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન તાપમાન (સે.ગ્રે)	33	34	33	34	34
લઘૃતમ તાપમાન (સે.ગ્રે)	20	20	21	21	21
વાદળની સ્થિતિ (ઓક્ટા)	1	0	0	0	1
મહત્વમાન ભેજ (%) સવાર	58	51	40	36	45
લઘૃતમ ભેજ (%) બપોરબાદ	32	29	26	21	20
પવનની ગતિ (કિમી/કલાક)	5	7	10	6	9
પવનની દિશા (ડિશ્રી)	157	87	38	26	342

૨. જામનગર જીલ્લો

તારીખ	દિવસ-૧ ૦૨-૧૧-૧૩	દિવસ-૨ ૦૩-૧૧-૧૩	દિવસ-૩ ૦૪-૧૧-૧૩	દિવસ-૪ ૦૫-૧૧-૧૩	દિવસ-૫ ૦૬-૧૧-૧૩
પેરામીટર					
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન (સે.ગ્રે)	30	31	30	31	31
લઘુત્તમ તાપમાન (સે.ગ્રે)	21	22	22	22	22
વાદળની સ્થિતિ (ઓક્ટા)	1	0	0	0	0
મહત્વમાન (%) સવાર	68	52	39	34	56
લઘુત્તમ ભેજ (%) બપોરબાદ	43	39	30	24	24
પવનની ગતિ (કિમી/કલાક)	7	10	12	8	11
પવનની દિશા (ડિગ્રી)	225	166	47	81	336

૩. અમરેલી જીલ્લો

તારીખ	દિવસ-૧ ૦૨-૧૧-૧૩	દિવસ-૨ ૦૩-૧૧-૧૩	દિવસ-૩ ૦૪-૧૧-૧૩	દિવસ-૪ ૦૫-૧૧-૧૩	દિવસ-૫ ૦૬-૧૧-૧૩
પેરામીટર					
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન (સે.ગ્રે)	32	33	32	33	33
લઘુત્તમ તાપમાન (સે.ગ્રે)	19	20	20	20	20
વાદળની સ્થિતિ (ઓક્ટા)	1	0	0	0	1
મહત્વમાન (%) સવાર	55	53	44	40	46
લઘુત્તમ ભેજ (%) બપોરબાદ	33	32	31	27	27
પવનની ગતિ (કિમી/કલાક)	5	8	11	7	7
પવનની દિશા (ડિગ્રી)	78	43	32	34	300

૪. સુરેન્દ્રનગર જીલ્લો

તારીખ	દિવસ-૧ ૦૨-૧૧-૧૩	દિવસ-૨ ૦૩-૧૧-૧૩	દિવસ-૩ ૦૪-૧૧-૧૩	દિવસ-૪ ૦૫-૧૧-૧૩	દિવસ-૫ ૦૬-૧૧-૧૩
પેરામીટર					
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન (સે.ગ્રે)	33	32	33	33	33
લઘુત્તમ તાપમાન (સે.ગ્રે)	22	23	23	23	22
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0
મહત્વમાન (%) સવાર	53	46	32	29	37
લઘુત્તમ ભેજ (%) બપોરબાદ	29	29	21	16	16
પવનની ગતિ (કિમી/કલાક)	4	9	11	5	7
પવનની દિશા (ડિગ્રી)	162	171	32	100	325

પત્રક-ક

ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ

હવામાન આગાહી

(તા. ૦૨-૧૧-૨૦૧૩ થી તા. ૦૬-૧૧-૨૦૧૩)

કૃષિ સલાહ:

તારીખ	દિવસ-૧ ૦૨-૧૧-૧૩	દિવસ-૨ ૦૩-૧૧-૧૩	દિવસ-૩ ૦૪-૧૧-૧૩	દિવસ-૪ ૦૫-૧૧-૧૩	દિવસ-૫ ૦૬-૧૧-૧૩
પેરામીટર					
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન તાપમાન (સે.ગ્રે)	33	34	33	34	34
લઘુત્વમાન તાપમાન (સે.ગ્રે)	20	20	21	21	21
વાદળી સ્થિતિ (ઓક્ટા)	1	0	0	0	1
મહત્વમાન ભેજ (%) સવાર	58	51	40	36	45
લઘુત્વમાન ભેજ (%) બપોરખાદ	32	29	26	21	20
પવનની ગતિ (કિમી/કલાક)	5	7	10	6	9
પવનની દિશા (ડિગ્રી)	157	87	38	26	342

સામાન્ય	<ul style="list-style-type: none"> ➤ ભારત સરકારશીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અતેના કૃષિ હવામાન વિભાગ, જૂનાગઢ કૃષિ યુનિવર્સિટી, તરફથી જાણવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના રાજકોટ જીલ્લામાં આગામી પાચ દિવસ તા. ૦૨-૧૧-૨૦૧૩ થી તા. ૦૬-૧૧-૨૦૧૩ દરમાન આકાશ ચોખ્યું રહેશે અને વરસાદની શક્યતા નથી. અને પવનની દિશા ઈશાન, અનિ અને પૂર્વ ની રહેવાની સંભાવના છે. પવનની ઝડપ ૫ થી ૧૦ કિમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્વમાન તાપમાન દિવસ દરમાન ઉત્ત થી ઉત્ત ડીગ્રી સેલ્ફિયસ જેટલું તેમજ લઘુત્વમાન રાત્રિ દરમાન ૨૦ થી ૨૧ ડિગ્રી સેલ્ફિયસ જેટલું રહેવાની સંભાવના છે. ➤ આ હવામાન પરિસ્થિતિમાં કપાસ અને શાકભાજી પાકોમાં ચુસીયા પ્રકારની જીવતોનો ઉપદ્રવ વધવાની શક્યતા હોય, ખેતરમાં ઊભા પાકમાં દેખરેખ રાખી જરૂરીયાત મુજબ જંતુનાશક દવા છાંટવી. ➤ જો ચોમાસુ પાક પૂર્ણ થય ગયેલ હોય તો લસણ, ચણા અને રાયડાનું પસંદગી મુજબ વાવેતર કરવું.
---------	--

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવત/જાત	કૃષિ સલાહ (૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
લસણ	વાવેતર	ગુજરાત લસણ-૨, ૪ અથવા સ્થાનિક	પાયામાં ૨૫ કિ.ગ્રા. નાઈટ્રોજન, ૫૦ કિ.ગ્રા. ફોસ્ફરસ અને ૫૦ કિ.ગ્રા. પોટાશ તેમજ ૨૫૦ કિ.ગ્રા. એરેનીનો ખોળ પ્રતિ હેક્ટર મુજબ આપી લસણની વાવણી કરવી.
ચણા	વાવેતર	ગુજરાત ચણા-૨ બીનપિયત માટે અને ગુજરાત ચણા-૩	પાયામાં ૨૦ કિ.ગ્રા. નાઈટ્રોજન અને ૪૦ કિ.ગ્રા. ફોસ્ફરસ પ્રતિ હેક્ટર મુજબ આપી ચણાની વાવણી કરવી. વાવતી વખતે ચણાના બીજાને ૨૫૦ ગ્રામ રાઈઝોબીયમ કલ્યરનો પ્રતિ ૧૦ કિ.ગ્રા. બીજ મુજબ ૫૮ પટ આપવો.
રાયડા	વાવેતર	રાય વરુણા અને ગુજરાત રાય-૩	પાયામાં ૩૦ કિ.ગ્રા. નાઈટ્રોજન, ૩૦ કિ.ગ્રા. ફોસ્ફરસ પ્રતિ હેક્ટર મુજબ આપી રાયમાની વાવણી કરવી.

કપાસ	જીડવા ખુલવા	લીલી પોપટી, શ્રીખસ, મોલોમશી  	નિયંત્રણ માટે ઈમીડાકલોપ્રીડ રો.૫ એસસી ૫ મીલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટવી.
		ખૂણિયા ટપકા, બળીયા ટપકા  	રોગ દેખાય કે તરત જ સ્ટ્રેપ્ટોસાયકલીન ૧ ગ્રામ + કોપર ઓક્ઝીકલોરાઈડ (૫૦ % વેટેબલ પાવડર) ૪૦ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્રણ કરી ૨ થી ૩ વખત છંટકાવ કરવો. કપાસના પાકમાં જડપી રીકવરી માટે સૂક્ષ્મ તત્વોયુક્ત NPK (૧૮-૧૮-૧૮) ૭૫ ગ્રામ ૧૦ લી પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
અરંડા	વૃદ્ધિ -કુલ	લીલી પોપટી, સર્ફદમાખી	તેના નિયંત્રણ માટે થાયોમીથોકઝામ ૨૫ ડબલ્યુઝી ૪ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટવી.
કુંગળી	વિકાસ	પાનનીટોયનોસુકારો /જાંબલી ઘાબાનો રોગ  શ્રીખસ	આ રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અને હેકજાકોનાજોલ ૧૫ મીલી દવાના ૧૦ લી પાણીમાં મિશ્ર કરી વારાફરતી બે છંટકાવ ૧૦ દિવસના અંતરે કરવા. તેના નિયંત્રણ માટે સ્પીનોસેડ ઉ મીલી ૧૦ લી પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
રીગણી અને ટમેટી	કુલ/ફળ	દુંખ અને – ફળ કોરી ખાનાર ઈયળ સર્ફદમાખી – તડતડીયા  	દુંખ અને ફળ કોરી ખાનાર ઈયળના નિયંત્રણ માટે નુકશાન થયેલ દુંખ અને ફળોને તોડીને દર અઠવારીએ નિયમીત અવશ્ય જમીનમાં દારી દેવા અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ પાણીમાં લેળવી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો સ્પીનોસેડ ૪૮ ઈસી અથવા રીનાક્ષીપીયર ઉ મીલી ૧૦ લી પાણીમાં મિશ્ર કરી દવા છાંટવી.
મરચી	કુલ/ફળ	શ્રીખસ  સુકારો	સ્પીનોસેડ ૪૮ ઉ મીલી અથવા એબામેક્ટીન ૧૦ મી.લી. દવા વારા ફરતી ૧૦ લીટર પાણીમાં મિશ્ર કરી ૧૦ દિવસના અંતરે છાંટવી. ૪૦ ગ્રામ ટ્રાયકોડર્મા કુગનો પાવડર ૧૦ લીટર પાણીમાં મિશ્ર કરી જમીનમાં થડની ફરતે મુળ વિસ્તારમાં નિતારવો.
વેલાવાળા શાકભાજી	કુલ/ફળ	તળધારો –	મેન્કોઝેબ ૨૭ ગ્રામ અથવા કલોરોયેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે નિયંત્રણ છંટકાવ કરવા.
સીતાકણ	ફળ	મીલીબગ/ચીકટો 	કવીનાલફોસ ૨૦ મીલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટવી.
પશુ/મરધા	પશુઓમાં ખરવા મોવાસાના રોગના નિયંત્રણ માટે રસીકરણ કરવું. તેમજ ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેશ્નીન અથવા એમીડાજ ૨ મી.લી. એક લીટર પાણીમાં નાખીને છાંટવી. જાનવરોની રહેઠાણની જગ્યાની વ્યવસ્થિત સફાઈ તેમજ માખીઓના ઉપદ્રવથી બચવા ફિનાઈલ છાંટવું.		