



હવામાન આગાહી આધારીત કૃષિ સલાહ બુલેટીન
ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ
તા. ૩૦-૦૪-૨૦૧૪ થી તા. ૦૪-૦૫-૨૦૧૪



(સંકલીત કૃષિ મૈસમ સલાહ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડીયા અને
ભારત મૈસમ વિભાગ દ્વારા સંપન્ન)

કૃષિ હવામાન બુલેટીન નં : ૬/૨૦૧૪

પાછલા અઠવાડીયાનું હવામાન

હવામાન પરિબળો	ઉત્તર સૌરાષ્ટ્ર
	રાજકોટ
વરસાદ (મી.મી)	0.0
મહત્વમાન તાપમાન (સે.ગ્રે)	39.5-43.5
લઘુત્વમાન તાપમાન (સે.ગ્રે)	23.0-26.0
ભેજ (%) સવાર	71-78
બપોરખાદ	17-23
પવનની ગતિ (કિમી/કલાક)	6.9-9.6

હવામાન આગાહી:

૧. રાજકોટ જીલ્લો

તારીખ	દિવસ-૧ ૩૦-૦૪-૧૪	દિવસ-૨ ૦૧-૦૫-૧૪	દિવસ-૩ ૦૨-૦૫-૧૪	દિવસ-૪ ૦૩-૦૫-૧૪	દિવસ-૫ ૦૪-૦૫-૧૪
પેરામીટર					
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન તાપમાન (સે.ગ્રે)	43	44	44	43	43
લઘુત્વમાન તાપમાન (સે.ગ્રે)	26	26	25	25	25
વાણની સ્થિતિ (ઓક્ટા)	0	0	0	0	0
મહત્વમાન ભેજ (%) સવાર	62	62	56	62	62
લઘુત્વમાન ભેજ (%) બપોરખાદ	22	21	25	31	32
પવનની ગતિ (કિમી/કલાક)	14	15	16	15	14
પવનની દિશા (રીશી)	290	270	270	260	270

૨. જામનગર જીવિતો

તારીખ	દિવસ-૧ ૩૦-૦૪-૧૪	દિવસ-૨ ૦૧-૦૫-૧૪	દિવસ-૩ ૦૨-૦૫-૧૪	દિવસ-૪ ૦૩-૦૫-૧૪	દિવસ-૫ ૦૪-૦૫-૧૪
પેરામીટર					
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન તાપમાન (સે.ગ્રે)	39	39	38	39	39
લઘુત્વમાન તાપમાન (સે.ગ્રે)	27	25	25	25	26
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0
મહત્વમાન ભેજ (%) સવાર	60	59	60	55	55
લઘુત્વમાન ભેજ (%) બપોરબાદ	35	35	41	47	48
પવનની ગતિ (કિમી/કલાક)	16	17	18	17	15
પવનની દિશા (ડીશ્રી)	280	260	260	250	270

૩. અમરેલી જીવિતો

તારીખ	દિવસ-૧ ૩૦-૦૪-૧૪	દિવસ-૨ ૦૧-૦૫-૧૪	દિવસ-૩ ૦૨-૦૫-૧૪	દિવસ-૪ ૦૩-૦૫-૧૪	દિવસ-૫ ૦૪-૦૫-૧૪
પેરામીટર					
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન તાપમાન (સે.ગ્રે)	42	42	43	41	41
લઘુત્વમાન તાપમાન (સે.ગ્રે)	27	25	25	25	25
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0
મહત્વમાન ભેજ (%) સવાર	57	64	68	65	65
લઘુત્વમાન ભેજ (%) બપોરબાદ	24	23	24	29	31
પવનની ગતિ (કિમી/કલાક)	15	14	14	13	14
પવનની દિશા (ડીશ્રી)	300	290	280	270	290

૪. સુરેન્દ્રનગર જીવિતો

તારીખ	દિવસ-૧ ૩૦-૦૪-૧૪	દિવસ-૨ ૦૧-૦૫-૧૪	દિવસ-૩ ૦૨-૦૫-૧૪	દિવસ-૪ ૦૩-૦૫-૧૪	દિવસ-૫ ૦૪-૦૫-૧૪
પેરામીટર					
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન તાપમાન (સે.ગ્રે)	41	40	39	37	38
લઘુત્વમાન તાપમાન (સે.ગ્રે)	27	26	26	26	26
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0
મહત્વમાન ભેજ (%) સવાર	50	62	66	65	64
લઘુત્વમાન ભેજ (%) બપોરબાદ	15	14	17	22	23
પવનની ગતિ (કિમી/કલાક)	17	16	16	15	15
પવનની દિશા (ડીશ્રી)	270	270	270	260	270

પત્રક-ક
ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ
હવામાન આગાહી
(તા. ૩૦-૦૪-૨૦૧૪ થી તા. ૦૪-૦૫-૨૦૧૪)

કુષી સલાહ :

સામાન્ય	> ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણ કુષી મૈસમ સેવા વિભાગ, જુનાગઢ કુષી યુનિવર્સિટી, તરઘડીયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના																																																						
	> રાજકોટ: જીલ્લામાં આગામી તા. ૩૦-૦૪-૨૦૧૪ થી તા. ૦૪-૦૫-૨૦૧૪ દરમયાન હવામાન સૂક્ષ્મ, સુર્ય પ્રકાશિત અને ચોખ્ખુ રહેશે. પવનની દિશા પણ્યમની રહેવાની અને પવનની ઝડપ ૧૪ થી ૧૬ ડિગ્રી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્વમાન દિવસ દરમયાન ૪૮ થી ૪૪ ડિગ્રી સેલ્ફીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રિ દરમયાન ૨૫ થી ૨૬ ડિગ્રી સેલ્ફીયસ જેટલું રહેવાની સંભાવના છે.																																																						
	<table border="1"> <thead> <tr> <th>તારીખ</th><th>દિવસ-૧ પેરામીટર ૩૦-૦૪-૧૪</th><th>દિવસ-૨ ૦૧-૦૫-૧૪</th><th>દિવસ-૩ ૦૨-૦૫-૧૪</th><th>દિવસ-૪ ૦૩-૦૫-૧૪</th><th>દિવસ-૫ ૦૪-૦૫-૧૪</th></tr> </thead> <tbody> <tr> <td>વરસાદ (મી.મી)</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr> <tr> <td>મહત્વમાન તાપમાન (સે.શે)</td><td>43</td><td>44</td><td>44</td><td>43</td><td>43</td></tr> <tr> <td>લઘુત્તમ તાપમાન (સે.શે)</td><td>26</td><td>26</td><td>25</td><td>25</td><td>25</td></tr> <tr> <td>વાદળની સ્થિતિ (ઓક્ટા)</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr> <tr> <td>મહત્વમાન ભેજ (%) સવાર</td><td>62</td><td>62</td><td>56</td><td>62</td><td>62</td></tr> <tr> <td>લઘુત્તમ ભેજ (%) બપોરબાદ</td><td>22</td><td>21</td><td>25</td><td>31</td><td>32</td></tr> <tr> <td>પવનની ગતિ (ડિગ્રી/કલાક)</td><td>14</td><td>15</td><td>16</td><td>15</td><td>14</td></tr> <tr> <td>પવનની દિશા (ડિગ્રી)</td><td>290</td><td>270</td><td>270</td><td>260</td><td>270</td></tr> </tbody> </table>	તારીખ	દિવસ-૧ પેરામીટર ૩૦-૦૪-૧૪	દિવસ-૨ ૦૧-૦૫-૧૪	દિવસ-૩ ૦૨-૦૫-૧૪	દિવસ-૪ ૦૩-૦૫-૧૪	દિવસ-૫ ૦૪-૦૫-૧૪	વરસાદ (મી.મી)	0	0	0	0	0	મહત્વમાન તાપમાન (સે.શે)	43	44	44	43	43	લઘુત્તમ તાપમાન (સે.શે)	26	26	25	25	25	વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0	મહત્વમાન ભેજ (%) સવાર	62	62	56	62	62	લઘુત્તમ ભેજ (%) બપોરબાદ	22	21	25	31	32	પવનની ગતિ (ડિગ્રી/કલાક)	14	15	16	15	14	પવનની દિશા (ડિગ્રી)	290	270	270	260	270
તારીખ	દિવસ-૧ પેરામીટર ૩૦-૦૪-૧૪	દિવસ-૨ ૦૧-૦૫-૧૪	દિવસ-૩ ૦૨-૦૫-૧૪	દિવસ-૪ ૦૩-૦૫-૧૪	દિવસ-૫ ૦૪-૦૫-૧૪																																																		
વરસાદ (મી.મી)	0	0	0	0	0																																																		
મહત્વમાન તાપમાન (સે.શે)	43	44	44	43	43																																																		
લઘુત્તમ તાપમાન (સે.શે)	26	26	25	25	25																																																		
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0																																																		
મહત્વમાન ભેજ (%) સવાર	62	62	56	62	62																																																		
લઘુત્તમ ભેજ (%) બપોરબાદ	22	21	25	31	32																																																		
પવનની ગતિ (ડિગ્રી/કલાક)	14	15	16	15	14																																																		
પવનની દિશા (ડિગ્રી)	290	270	270	260	270																																																		
	> જામનગર: જીલ્લામાં આગામી તા. ૩૦-૦૪-૨૦૧૪ થી તા. ૦૪-૦૫-૨૦૧૪ દરમયાન હવામાન સૂક્ષ્મ, સુર્ય પ્રકાશિત અને ચોખ્ખુ રહેશે. પવનની દિશા પણ્યમની રહેવાની અને પવનની ઝડપ ૧૫ થી ૧૮ ડિગ્રી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્વમાન દિવસ દરમયાન ૩૮ થી ૩૮ ડિગ્રી સેલ્ફીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રિ દરમયાન ૨૫ થી ૨૬ ડિગ્રી સેલ્ફીયસ જેટલું રહેવાની સંભાવના છે.																																																						
	<table border="1"> <thead> <tr> <th>તારીખ</th><th>દિવસ-૧ પેરામીટર ૩૦-૦૪-૧૪</th><th>દિવસ-૨ ૦૧-૦૫-૧૪</th><th>દિવસ-૩ ૦૨-૦૫-૧૪</th><th>દિવસ-૪ ૦૩-૦૫-૧૪</th><th>દિવસ-૫ ૦૪-૦૫-૧૪</th></tr> </thead> <tbody> <tr> <td>વરસાદ (મી.મી)</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr> <tr> <td>મહત્વમાન તાપમાન (સે.શે)</td><td>39</td><td>39</td><td>38</td><td>39</td><td>39</td></tr> <tr> <td>લઘુત્તમ તાપમાન (સે.શે)</td><td>27</td><td>25</td><td>25</td><td>25</td><td>26</td></tr> <tr> <td>વાદળની સ્થિતિ (ઓક્ટા)</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr> <tr> <td>મહત્વમાન ભેજ (%) સવાર</td><td>60</td><td>59</td><td>60</td><td>55</td><td>55</td></tr> <tr> <td>લઘુત્તમ ભેજ (%) બપોરબાદ</td><td>35</td><td>35</td><td>41</td><td>47</td><td>48</td></tr> <tr> <td>પવનની ગતિ (ડિગ્રી/કલાક)</td><td>16</td><td>17</td><td>18</td><td>17</td><td>15</td></tr> <tr> <td>પવનની દિશા (ડિગ્રી)</td><td>280</td><td>260</td><td>260</td><td>250</td><td>270</td></tr> </tbody> </table>	તારીખ	દિવસ-૧ પેરામીટર ૩૦-૦૪-૧૪	દિવસ-૨ ૦૧-૦૫-૧૪	દિવસ-૩ ૦૨-૦૫-૧૪	દિવસ-૪ ૦૩-૦૫-૧૪	દિવસ-૫ ૦૪-૦૫-૧૪	વરસાદ (મી.મી)	0	0	0	0	0	મહત્વમાન તાપમાન (સે.શે)	39	39	38	39	39	લઘુત્તમ તાપમાન (સે.શે)	27	25	25	25	26	વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0	મહત્વમાન ભેજ (%) સવાર	60	59	60	55	55	લઘુત્તમ ભેજ (%) બપોરબાદ	35	35	41	47	48	પવનની ગતિ (ડિગ્રી/કલાક)	16	17	18	17	15	પવનની દિશા (ડિગ્રી)	280	260	260	250	270
તારીખ	દિવસ-૧ પેરામીટર ૩૦-૦૪-૧૪	દિવસ-૨ ૦૧-૦૫-૧૪	દિવસ-૩ ૦૨-૦૫-૧૪	દિવસ-૪ ૦૩-૦૫-૧૪	દિવસ-૫ ૦૪-૦૫-૧૪																																																		
વરસાદ (મી.મી)	0	0	0	0	0																																																		
મહત્વમાન તાપમાન (સે.શે)	39	39	38	39	39																																																		
લઘુત્તમ તાપમાન (સે.શે)	27	25	25	25	26																																																		
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0																																																		
મહત્વમાન ભેજ (%) સવાર	60	59	60	55	55																																																		
લઘુત્તમ ભેજ (%) બપોરબાદ	35	35	41	47	48																																																		
પવનની ગતિ (ડિગ્રી/કલાક)	16	17	18	17	15																																																		
પવનની દિશા (ડિગ્રી)	280	260	260	250	270																																																		
	> અમરેલી: જીલ્લામાં આગામી તા. ૩૦-૦૪-૨૦૧૪ થી તા. ૦૪-૦૫-૨૦૧૪ દરમયાન હવામાન સૂક્ષ્મ, સુર્ય પ્રકાશિત અને અંશતઃ વાદળધાયુ રહેશે. પવનની દિશા વાયવ્ય અને પણ્યમની રહેવાની અને પવનની ઝડપ ૧૩ થી ૧૫ ડિગ્રી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્વમાન દિવસ દરમયાન ૪૧ થી ૪૩ ડિગ્રી સેલ્ફીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રિ દરમયાન ૨૫ થી ૨૭ ડિગ્રી સેલ્ફીયસ જેટલું રહેવાની સંભાવના છે.																																																						

તારીખ	દિવસ-૧ ૩૦-૦૪-૧૪	દિવસ-૨ ૦૧-૦૫-૧૪	દિવસ-૩ ૦૨-૦૫-૧૪	દિવસ-૪ ૦૩-૦૫-૧૪	દિવસ-૫ ૦૪-૦૫-૧૪
પેરામીટર					
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન (સે.ગ્રે)	42	42	43	41	41
લઘુત્તમ તાપમાન (સે.ગ્રે)	27	25	25	25	25
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0
મહત્વમાન (%) સવાર	57	64	68	65	65
લઘુત્તમ ભેજ (%) બપોરબાદ	24	23	24	29	31
પવનની ગતિ (કિમી/કલાક)	15	14	14	13	14
પવનની દિશા (ડિગ્રી)	300	290	280	270	290

➤ સુરેન્દ્રગર જીવલામાં આગામી તા. ૩૦-૦૪-૨૦૧૪ થી તા. ૦૪-૦૫-૨૦૧૪ દરમયાન હવામાન સૂક્ક, સુર્ય પ્રકાશિત અને અંશતઃ વાદળાયુ રહેશે. પવનની દિશા વાયવ્ય અને પણ્યમની રહેવાની અને પવનની ઝડપ ૧૫ થી ૧૮ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્વમાન દિવસ દરમયાન ૩૮ થી ૪૧ ડિગ્રી સેલ્ફીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રિ દરમયાન ૨૬ થી ૨૮ ડિગ્રી સેલ્ફીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧ ૩૦-૦૪-૧૪	દિવસ-૨ ૦૧-૦૫-૧૪	દિવસ-૩ ૦૨-૦૫-૧૪	દિવસ-૪ ૦૩-૦૫-૧૪	દિવસ-૫ ૦૪-૦૫-૧૪
પેરામીટર					
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન (સે.ગ્રે)	41	40	39	37	38
લઘુત્તમ તાપમાન (સે.ગ્રે)	27	26	26	26	26
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0
મહત્વમાન (%) સવાર	50	62	66	65	64
લઘુત્તમ ભેજ (%) બપોરબાદ	15	14	17	22	23
પવનની ગતિ (કિમી/કલાક)	17	16	16	15	15
પવનની દિશા (ડિગ્રી)	270	270	270	260	270

- ગરમ લું પડવાની સંભાવના હોય, ઉનાળું વાવેતર કરેલ પાકોને જરૂરીયાત મુજબ પિયત આપવું.
 ➤ હાલ ઉનાળું મગફળી અને તલના પાકમાં ગરમ અને ભેજવાળા વાતરવરણના કારણે પાન કથીરીનો ઉપદ્રવ થવાની શક્યતા હોય પાકનું સતત નીરીક્ષણ કરતા રહેવું. જો ઉપદ્રવ જોવા મળે તો પ્રોપરગાઈટ ૧૦ મીલી ૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકવ કરવો.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટર પાણીમાં મીશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
ખાતરની ખાડ	પાકું ખાતર પલટાવવું	—	<p>ઢોરના મળમુંન અને ખેત કચરામાંથી પૂર્ણ સરી ગયેલ ખાતરને ખાડમાંથી બહાર કાઢવું.</p> 

ઉનાળુ મગફળી	કુલ, સુયા બેસવા, અને પોપટા બંધાવા	-	<ul style="list-style-type: none"> ➢ કુલ અને સુયા અવસ્થાએ હળવુ પિયત આપવુ. 
ઉનાળુ તલ	વૃધિ/કુલ/બેઈટા બેસવા	સફેદ માખી	<ul style="list-style-type: none"> ➢ તેના નિયંત્રણ માટે લીમડાનું તેલ ૫૦ મીલી અથવા ટ્રાયાફોસ ૪૦ ટકા ૨૦ મીલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકવ કરવો. ➢ તેમજ તલના પાકને કુલ અવસ્થાએ હળવુ પિયત આપવુ. ત્યારબાદ જમીનની પ્રત સુજલ ૧૦-૧૨ ટિવસના ગાળે પિયત આપવુ. વધારે પ્રમાણમાં ઢૂંકા અંતરે પિયત આપવા નહિ. 
ઉનાળુ મગ	કુલ અને શીંગ	લીલી ઈયણ	<ul style="list-style-type: none"> ➢ લીલોળીનું તેલ ૫૦ મીલી અથવા લીલોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો ઈન્ડોક્સાકાર્બ ૧૦ મીલી /૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકવ કરવો. 
ઉનાળુ ગમ ગુવાર	કુલ અને શીંગ	શ્રીપ્સ, સફેદ માખી	<ul style="list-style-type: none"> ➢ તેના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા ૩ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકવ કરવો. ➢ જો ભુકીછારાનો ઉપદ્રવ જણાય તો હેકઝાકોનાજોલ ૫ ઈસી/ ૧૦ મીલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકવ કરવો.

કપાસ અને ઘઉં	કરાંઠી ઉપાડવી કાપણી	 	<p>કપાસની કરાંઠીને રોટાવેટર/ચકરડી(ચોપર)ની મદદથી ટૂકડા કરીને જમીનમાં ભેણવી/દાટી દેવી જેથી જમીનમાં સેન્ટ્રીય તત્વ ઉમેરાવાથી જમીનના સ્વાસ્થ્યમાં સુધારો થશે.</p> <p>ઘઉની કાપણી હાર્ટેસ્ટરથી કર્યા પછી ખેતરમાં રહેલ ઘઉના ભાડા-પરાળ સળગાવતું નહીં. પરંતુ તેને થોડા ટિવસ સુર્યના તડકામાં તપવા દેવા અને ત્યારબાદ બપોરબાદ સમાર, ટ્રેક્ટરનું સુપડુ કે રોટાવેટરથી ભૂકો કરી જમીનમાં ભેણવી દેવા.</p>
ચારાની જીવાર	વૃદ્ધિ	મોલો અને કુંપળની ઈયળ	<p>તેના નિયંત્રણ માટે અનુકૂમે એસીટેમાપીડ 20 ટકા ઉ ગ્રામ અને લેમા સાયદેલોશ્રીન ૫ ટકા ૧૦ મી.લી./૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકાવ કરવો. દવા છંટકાવ બાદ ૨૦ ટિવસ સુધી પશુઓને ખવડાવવી નહીં.</p>  
રીગણી અને ટ્યેટી	કુલ/ફળ	સફેદ માખી દૂંખ અને ફળ કોરી ખાનાર ઈયળ	<p>લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટેમાપીડ 20 ટકા ઉ ગ્રામ /૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકાવ કરવો.</p> <p>દૂંખ અને ફળ કોરી ખાનાર ઈયળના નિયંત્રણ માટે નુકશાન થયેલ દૂંખ અને ફળોને તોડીને દર અઠવાડીયે નિયમીત જમીનમાં દાટી દેવા અને જો વધુ ઉપદ્રવ હોય તો સ્પીનોસેડ ૪૮ ઈસી ઉ મીલી /૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકાવ કરવો. દવા છંટકાવ બાદ ૮ ટિવસ સુધી ફળ ઉતારવા નહીં.</p>
મરચી	કુલ/ફળ	શ્રીપસ	<p>પ્રોફેનોકોસ ૨૦ મી.લી. અને લીબોળીનું તેલ ૫૦ મીલી/૧૦ લી પાણીમાં મિશ્ર કરી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ કરી છંટકાવ કરવો.</p> 
ભાડા	કુલ/ફળ	ચુસીયા તથા દૂંખ અને ફળ કોરી ખાનાર ઈયળ –	<p>દૂંખ અને ફળ કોરી ખાનાર ઈયળના નિયંત્રણ માટે નુકશાન થયેલ દૂંખ અને ફળોને તોડીને જમીનમાં દાટી દેવા અને લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ અથવા લીબોળીનું તેલ ૫૦ મીલી/૧૦ લી પાણીમાં મિશ્ર નો છંટકાવ ૧૫ ટિવસના અંતરે ચાર</p>

			વખત કરવો.
ગુવાર	વૃદ્ધિ/કુલ /શીંગ	શ્રીપસ, સર્કેદ માધી	<p>➢ તેના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા પ ગ્રામ અને લીબોળીનું તેલ ૫૦ મીલી /૧૦ લીટર પાણીમાં મીશ્ર કરી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણનો છંટકવ કરવો.</p>
વેલાવાળા શાકભાજી	વૃદ્ધિ/ કુલ	પાન કોરીયુ , શ્રીપસ, સર્કેદ માધી – તળાછારો –	<p>➢ તેના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા પ ગ્રામ અને લીબોળીનું તેલ ૫૦ મીલી /૧૦ લીટર પાણીમાં મીશ્ર કરી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણનો છંટકવ કરવો.</p> <p>મેન્કોરેલ ૨૭ ગ્રામ અથવા કલોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે નષ્ટ છંટકવ કરવા.</p> 
અંબો	ફળ વિકાસ	ફળમાની પિયત	<p>તેના નિયંત્રણ માટે મીથાઈલ યુજીનોલ ટ્રેપનો ઉપયોગ કરી ફળમાનીનું નિયંત્રણ કરવું.</p> <p>અંબાના પાકમાં હળવુ પિયત આપવુ.</p> 
વીળુ	ફળનો વિકાસ	સાયલા અને શ્રીપસ	<p>તેના નિયંત્રણ માટે ઈમીડાકલોપ્રીડ ૧૭.૮ ટકા પ મી.લી. /૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકવ કરવો.</p>   
પણુ/મરવા	તાપમાન વધારે રહેવાના કારણે ઠોરને દિવસના સમયે છાંયડામાં રાખવા અને પીવાના સ્વર્ચ પાણીની વ્યવસ્થા કરવી.		<p>પણુઓમાં ગળસુંઢો તથા ગાંઠીયા તાવના નિયંત્રણ માટે રસીકરણ કરવુ અને કરમીયાની ગોળી આપવી. દુધાળા પણુના આઉને જીક ઓક્સાઈડ અથવા બોરીક પાવડર થી બરાબર સાફ કરવુ. તેમજ ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથીન</p> 

અથવા એમીટાજ ર મી.લી. એક લીટર પાણીમાં નાખીને છાંટવી. દુધાળા પશુઓને ઉનાળામાં ઠડક આપવા પાણીનો છંટકાવ કરવો. બપોરના સમયે મરવાઓનું પરિવહન ટાળવું. મરવા ઘર ની પણ્યમ દિશામાં પડદા લગાવવા અને તેમની ઉપર બપોરના સમયે પાણીનો છંટકાવ કરવો.

જા.નં. જુક્કુયુ/ સંવૈ(સુખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૧૪
તરધીયા(રાજકોટ). તારીખ: ૨૮-૦૪-૨૦૧૪

પ્રતિ,
તંત્રીશ્રી/ ડાયરેક્ટરશ્રી,
રાજકોટ.
ઉપરોક્ત માહિતી આપના દેનિક સમાચાર પત્રોમાં, રેડિયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિનામૂલ્યે પ્રસિદ્ધ કરવા નમ્ર વિનંતી છે. સહકારની અપેક્ષા સહ આભાર.

સંશોધન વૈજ્ઞાનિક (સૂકી ખેતી)
જુનાગઢ કૃષિ યુનિવર્સિટી
તરધીયા



**Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523**



[Agro met Advisory Service bulletin no.- 9 for the Rajkot district](#)
(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
(Period 30th April To 4th May - 2014)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	0.0
Maximum temperature (°C)	39.5-43.5
Minimum temperature (°C)	23.0-26.0
Relative Humidity (%) I	71-78
Relative Humidity (%) II	17-23
Wind speed (Kmph)	6.9-9.6


Weather forecast until 08-30 hrs of 04/05/2014

(1) RAJKOT DISTRICT:

	30/04/14	01/05/14	02/05/14	03/05/14	04/05/14
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	43	44	44	43	43
Min.Temp.(°C)	26	26	25	25	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	62	62	56	62	62
RH-II (%)	22	21	25	31	32
Wind Speed (kmph)	14	15	16	15	14
Wind Direction(deg.)	290	270	270	260	270

Agro meteorological Advisories:

General Advice	<p>➤ Rain fall: Dry, sunny, hot and clear weather forecasted for next five days.</p> <p>➤ Temperature: Maxmun and minimum temperature likely to be in range of 43-44 and 25-26°C, respectively.</p> <p>➤ Relative Humidity: The RH is likely to be in the range of 21-62 %</p> <p>➤ Wind Direction and Speed: Direction and speed of wind is likely to be W and 14-16 km/h, respectively.</p> <p>➤ Rise in maximum temperature by 2-3 °c is forecasted for next five days, leading to development of heat wave condition, irrigate summer crops as per need.</p>		
Crops	Stage	Insect/Pest/Variety	Advisory
Farm compost	Turning of	-	Dig out completely composted material made from animal dung, urine, litter and farm waste.

	composted material		
Summer groundnut	Flowering pegging/ Pod development	White fly and prodenia	Apply light irrigation at flowering & pegging stage 
Summer Sesame	Flowering /capsule formation	White fly	Under present weather condition infestation of whiteflies observed in summer sown sesame. To control them spray acetamiprid 20 % SP@ 5.0 gm /10 lit of water. 
Green gram	Flowering / Pod formation	Heliothis	Under present weather condition severe attack of Heliothis observed in summer sown green gram. To control heliothis spray neem seed extract 5 % or neem seed oil 50 ml and quinalphos 20 ml / 10 lit of water. In case of heavy attack make immediate spray of Indoxacarb 10 ml/ 10 liter of water. 
Gum Guar	Flowering / pod	Powdery mildew	To control PM spray Hexaconazole 5 % EC 15 ml/10 lit of water
Cotton &	pulling of stalk	 Mobile chopper	Do in situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator for improving soil health.

Wheat	Harvested		Do not burn the wheat straw, stubbles and other left on materials after harvesting of wheat crop by wheat harvester, but mix it in soil through planking, rotavator and cultivator.
Fodder sorghum	Growth	Ahid and shoot borer	Spray acetamiprid 20 % SP@ 5.0 gm and <i>lambda-cyhalothrin</i> 5 % EC, 10 ml/10 litre of water, respectively.   
Lady finger	Flowering /fruiting	Sucking pest thrips, white fly & Shoot and fruit borer	To control them spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Collect damaged fruits and buried deep in soil.
Ridge gourd/ Bottle gourd/ Sponge gourd / Bitter gourd	Flowering	Leaf minor, Thrips, white fly Downy mildew & Alternaria blight 	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Cluster bean	Flowering /pod	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
Brinjal	Flowering /fruiting	Sucking pest & Fruit borer	To control them spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Collect damaged fruits and buried deep in soil.
Chili	Flowering & fruiting	Thrips → 	In present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofosin 10 litres of water. 
Mango	Fruiting	Blossom midge and Leaf hopper → Fruit fly	To prevent damage spray imidacloprid 5 ml/10 litres of water. Spray mixture of Fenthion 10 ml or DDVP 4 ml

		Irrigation	and Methyle ugenol 10 ml in 10 litres of water Apply light irrigation 
Lemon	Fruiting	Psylla and thrips 	To prevent damage spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.   
Livestock & poultry 			Under present weather condition shower the animals with cold water two to three times in a day. Increase the green fodder quantity for feeding. Supplimentry feeding of vitamins A and D should apply. Do Vaccination for Haemorrhageic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water. Put curtains on west side of poultry house and sprinkle water on curtains during noon time.

GKMS Unit, DFRS, JAU, Targhadia



**Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523**



Agro met Advisory Service bulletin for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 30th April To 4th May - 2014)

Weather forecast until 08-30 hrs of 04/05/2014

(2) JAMNAGAR DISTRICT:

	30/04/14	01/05/14	02/05/14	03/05/14	04/05/14
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	39	39	38	39	39
Min.Temp.(°C)	27	25	25	25	26
Total CC (octa)	0	0	0	0	0
RH-I (%)	60	59	60	55	55
RH-II (%)	35	35	41	47	48
Wind Speed (kmph)	16	17	18	17	15
Wind Direction(deg.)	280	260	260	250	270

Agro meteorological Advisories:-

General Advice	<p>➤ Rain fall: Dry, sunny and clear weather forecasted for next five.</p> <p>➤ Temperature: Maxmun and minimum temperature likely to be in range of 38-39 and 25-27°C, respectively.</p> <p>➤ Relative Humidity: The RH is likely to be in the range of 35-60 %</p> <p>➤ Wind Direction and Speed: Direction and speed of wind is likely to be W and 15-18 km/h, respectivley.</p> <p>➤ Rise in maximum temperature by 1-2 °c forecasted for next five days, so irrigate summer crops as per need.</p>			
Crops	Stage	Insect/Pest/Variety	Advisory	
Farm compost	Turning of composted material	-	Dig out completely composted material made from animal dung, urine, litter and farm waste. 	
Summer groundnut	Flowering pegging/ Pod development	White fly and prodenia	Apply light irrigation at flowering & pegging stage 	

Summer Sesame	Flowering /capsule formation	White fly 	Under present weather condition infestation of whiteflies observed in summer sown sesame. To control them spray acetamiprid 20 % SP@ 5.0 gm/10 lit of water.  
Green gram	Flowering / Pod formation	Heliothis 	Under present weather condition severe attack of Heliothis observed in summer sown green gram. To control heliothis spray neem seed extract 5 % or neem seed oil 50 ml and quinalphos 20 ml / 10 lit of water. In case of heavy attack make immediate spray of Indoxacarb 10 ml/ 10 liter of water. 
Gum Guar	Flowering / pod	Powdery mildew	To control PM spray Hexaconazole 5 % EC 15 ml/10 lit of water
Cotton & Wheat	pulling of stalk Harvested	 	Do in situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator for improving soil health. Do not burn the wheat straw, stubbles and other left on materials after harvesting of wheat crop by wheat harvester, but mix it in soil through planking, rotavator and cultivator.
Fodder sorghum	Growth	Ahid and shoot borer	Spray acetamiprid 20 % SP@ 5.0 gm and lambda-cyhalothrin 5 % EC, 10 ml/10 litre of water, respectively.   

Lady finger	Flowering /fruiting	Sucking pest thrips, white fly & Shoot and fruit borer	To control them spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Collect damaged fruits and buried deep in soil.
Ridge gourd/ Bottle gourd/ Sponge gourd / Bitter gourd	Flowering	Leaf minor, Thrips, white fly Downy mildew & Alternaria blight 	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Cluster bean	Flowering /pod	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
Brinjal	Flowering /fruiting	Sucking pest & Fruit borer	To control them spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Collect damaged fruits and buried deep in soil.
Chili	Flowering & fruiting	Thrips → 	In present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofos in 10 litres of water. 
Mango	Fruiting	Blossom midge and Leaf hopper → Fruit fly Irrigation	To prevent damage spray imidacloprid 5 ml/10 litres of water. Spray mixture of Fenthion 10 ml or DDVP 4 ml and Methyle ugenol 10 ml in 10 litres of water Apply light irrigation 

Lemon	Fruiting	Psylla and thrips 	To prevent damage spray imidacloprid 30.5 % 5 ml / 10 lit of water . 
Livestock & poultry 	Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water. Shower the animals with cold water two to three times in a day. Increase the green fodder quantity for feeding. Supplimentry feeding of vitamins A and D should apply. Put curtains on west side of poultry house and sprinkle water on curtains during noon time.		

GKMS Unit, DFRS, JAU, Targhadia



**Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523**



Agro met Advisory Service bulletin for the Amreli district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 30th April To 4th May - 2014)

Weather forecast until 08-30 hrs of 04/05/2014

(3) AMRELI DISTRICT:

	30/04/14	01/05/14	02/05/14	03/05/14	04/05/14
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	42	42	43	41	41
Min.Temp.(°C)	27	25	25	25	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	57	64	68	65	65
RH-II (%)	24	23	24	29	31
Wind Speed (kmph)	15	14	14	13	14
Wind Direction(deg.)	300	290	280	270	290

Agro meteorological Advisories :

General Advice	<p>➤ Rain fall: Dry, sunny, hot and clear weather forecasted for next five days.</p> <p>➤ Temperature: Maximum and minimum temperature likely to be in range of 41-43 and 25-27°C, respectively.</p> <p>➤ Relative Humidity: The RH is likely to be in the range of 23-68 %</p> <p>➤ Wind Direction and Speed: Direction and speed of wind is likely to be NW and 13-14 km/h, respectively.</p> <p>➤ Rise in maximum temperature by 1-2 °c forecasted for next five days, so irrigate summer crops as per need.</p>
-----------------------	--

Crops	Stage	Insect/Pest/Variety	Advisory
Farm compost	Turning of composted material	-	Dig out completely composted material made from animal dung, urine, litter and farm waste. 
Summer groundnut	Flowering pegging/ Pod development	White fly and prodenia	Apply light irrigation at flowering & pegging stage

			 	
Summer Sesame	Flowering /capsule formation	White fly		Under present weather condition infestation of whiteflies observed in summer sown sesame. To control them spray acetamiprid 20 % SP@ 5.0 gm /10 lit of water.  
Green gram	Flowering / Pod formation	Heliothis		Under present weather condition severe attack of Heliothis observed in summer sown green gram. To control heliothis spray neem seed extract 5 % or neem seed oil 50 ml and quinalphos 20 ml / 10 lit of water. In case of heavy attack make immediate spray of Indoxacarb 10 ml/ 10 liter of water. 
Gum Guar	Flowering / pod	Powdery mildew		To control PM spray Hexaconazole 5 % EC 15 ml/10 lit of water
Cotton & Wheat	pulling of stalk Harvested	Mobile chopper	 	Do in situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator for improving soil health. Do not burn the wheat straw, stubbles and other left on materials after harvesting of wheat crop by wheat harvester, but mix it in soil through planking, rotavator and cultivator.
Fodder sorghum	Growth	Ahid and shoot borer		Spray acetamiprid 20 % SP@ 5.0 gm and lambda-cyhalothrin 5 % EC , 10 ml/10 litre of water, respectively.

			
Lady finger	Flowering /fruiting	Sucking pest thrips, white fly & Shoot and fruit borer	To control them spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Collect damaged fruits and buried deep in soil.
Ridge gourd/ Bottle gourd/ Sponge gourd / Bitter gourd	Flowering	Leaf minor, Thrips, white fly Downy mildew & Alternaria blight 	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Cluster bean	Flowering /pod	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
Brinjal	Flowering /fruiting	Sucking pest & Fruit borer	To control them spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Collect damaged fruits and buried deep in soil.
Chili	Flowering & fruiting	Thrips → 	In present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofos in 10 litres of water. 
Mango	Fruiting	Blossom midge and Leaf hopper → Fruit fly Irrigation	To prevent damage spray imidacloprid 5 ml/10 litres of water. Spray mixture of Fenthion 10 ml or DDVP 4 ml and Methyle ugenol 10 ml in 10 litres of water Apply light irrigation

				
Lemon	Fruiting	Psylla and thrips 	To prevent damage spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.	  
Livestock & poultry 	Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water. Shower the animals with cold water two to three times in a day. Increase the green fodder quantity for feeding. Supplimentry feeding of vitamins A and D should apply. Put curtains on west side of poultry house and sprinkle water on curtains during noon time.			

GKMS Unit, DFRS, JAU, Targhadia



**Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523**



Agro met Advisory Service bulletin for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 30th April To 4th May - 2014)

Weather forecast until 08-30 hrs of 04/05/2014

(4) SURENDRANAGAR DISTRICT:

	30/04/14	01/05/14	02/05/14	03/05/14	04/05/14
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	41	40	39	37	38
Min.Temp.(°C)	27	26	26	26	26
Total CC (octa)	0	0	0	0	0
RH-I (%)	50	62	66	65	64
RH-II (%)	15	14	17	22	23
Wind Speed (kmph)	17	16	16	15	15
Wind Direction(deg.)	270	270	270	260	270

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none">➤ Rain fall: Dry, sunny and clear weather forecasted for next five days.➤ Temperature: Maxmun and minimum temperature likely to be in range of 37-41°C and 26-27°C, respectively.➤ Relative Humidity: The RH is likely to be in the range of 14-66 %➤ Wind Direction and Speed: Direction and speed of wind is likely to W and 14-17 km/h, respectively.➤ Under prevailing weather irrigate summer crops as per need.
-----------------------	--

Crops	Stage	Insect/Pest/Variety	Advisory
Farm compost	Turning of composted material	-	Dig out completely composted material made from animal dung, urine, litter and farm waste. 
Summer groundnut	Flowering pegging/ Pod development	White fly and prodenia	Apply light irrigation at flowering & pegging stage 

Summer Sesame	Flowering/capsule formation	White fly 	Under present weather condition infestation of whiteflies observed in summer sown sesame. To control them spray acetamiprid 20 % SP@ 5.0 gm /10 lit of water.  
Green gram	Flowering/ Pod formation	Heliothis 	Under present weather condition severe attack of Heliothis observed in summer sown green gram. To control heliothis spray neem seed extract 5 % or neem seed oil 50 ml and quinalphos 20 ml / 10 lit of water. In case of heavy attack make immediate spray of Indoxacarb 10 ml/ 10 liter of water. 
Gum Guar	Flowering/ pod	Powdery mildew	To control PM spray Hexaconazole 5 % EC 15 ml/10 lit of water
Cotton & Wheat	pulling of stalk Harvested	Mobile chopper 	Do in situ chopping & burying of cotton stalk in soil by mobile chopper and rotavator for improving soil health. Do not burn the wheat straw, stubbles and other left on materials after harvesting of wheat crop by wheat harvester, but mix it in soil through planking, rotavator and cultivator.
Fodder sorghum	Growth	Ahid and shoot borer	Spray acetamiprid 20 % SP@ 5.0 gm and lambda-cyhalothrin 5 % EC , 10 ml/10 litre of water, respectively.   

Lady finger	Flowering /fruiting	Sucking pest thrips, white fly & Shoot and fruit borer	To control them spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Collect damaged fruits and buried deep in soil.
Ridge gourd/ Bottle gourd/ Sponge gourd / Bitter gourd	Flowering	Leaf minor, Thrips, white fly Downy mildew & Alternaria blight 	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Cluster bean	Flowering/pod	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
Brinjal	Flowering/fruiting	Sucking pest & Fruit borer	To control them spray neem seed extract 5 % or neem seed oil 50 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Collect damaged fruits and buried deep in soil.
Chili	Flowering & fruiting	Thrips → 	In present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofosin 10 litres of water. 
Mango	Fruiting	Blossom midge and Leaf hopper → Fruit fly Irrigation	To prevent damage spray imidaclorpid 5 ml/10 litres of water. Spray mixture of Fenthion 10 ml or DDVP 4 ml and Methyle ugenol 10 ml in 10 litres of water Apply light irrigation 

Lemon	Fruiting	Psylla and thrips	To prevent damage spray imidacloprid 30.5 % 5 ml / 10 lit of water ml/10 lit of water.
		   	

Livestock & poultry



Do Vaccination for **Haemorrhagic septicemia** (H.S.) disease and **Blue Quarter** (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water. Shower the animals with cold water two to three times in a day. Increase the green fodder quantity for feeding. Supplimentary feeding of vitamins A and D should apply. Put curtains on west side of poultry house and sprinkle water on curtains during noon time.

GKMS Unit, DFRS, JAU, Targhadia