



ગ્રામીણ કૃષિ મૌસમ સેવા

હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન

ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ

તા.૦૮-૦૪-૨૦૨૦ થી તા.૧૨-૦૪-૨૦૨૦



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કે.યુ. તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિ હવામાન બુલેટીન નં. ૩/૨૦૨૦

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

હવામાન પરિબળો	
વરસાદ(મી.મી)	0.0
મહત્તમ તાપમાન(સે.ગ્રે)	37.0-40.0
લઘુતમ તાપમાન(સે.ગ્રે)	19.0-22.0
ભેજ (%)સવાર	61-74
ભેજ (%)બપોરબાદ	20-25
પવનની ગતિ(કિમી/કલાક)	3.9-6.1

હવામાન આગાહી :પત્રક-ક

ઉત્તરસૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.૦૮-૦૪-૨૦૨૦ થી ૧૨-૦૪-૨૦૨૦

કૃષિ સલાહ :

સામાન્ય	<p>➤ ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણકૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના</p> <p>રાજકોટ : જીલ્લામાં આગામી તા.૦૮-૦૪-૨૦૨૦ થી તા.૧૨-૦૪-૨૦૨૦ દરમિયાન હવામાન સૂકું, ગરમ અને યોખ્ખું રહેશે. પવનની દિશા પશ્ચિમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૯ થી ૧૪ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૯-૪૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમિયાન ૨૨-૨૩ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.</p>					
	તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
	પેરામીટર	૦૮-૦૪-૨૦	૦૯-૦૪-૨૦	૧૦-૦૪-૨૦	૧૧-૦૪-૨૦	૧૨-૦૪-૨૦
	વરસાદ(મી.મી.)	0	0	0	0	0
	મહત્તમતાપમાન(સે.ગ્રે)	39	39	39	40	41
	લઘુતમતાપમાન(સે.ગ્રે)	22	22	22	22	23
	વાદળનીસ્થિતિ(ઓકટા)	0	0	0	0	1
	મહત્તમભેજ (%)સવાર	80	78	85	76	55
	લઘુતમભેજ (%) બપોરબાદ	25	34	28	26	23
	પવનનીગતિ(કિમી/કલાક)	14	13	11	9	12
	પવનનીદિશા(ડીગ્રી)	255	266	280	308	320
	<p>મોરબી : જીલ્લામાં આગામી તા.૦૮-૦૪-૨૦૨૦ થી તા.૧૨-૦૪-૨૦૨૦ દરમિયાન હવામાન સૂકું, ગરમ અને યોખ્ખું રહેશે. પવનની દિશા પશ્ચિમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૯ થી ૧૪ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૯-૪૧ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમિયાન ૨૨-૨૩ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.</p>					
	તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
	પેરામીટર	૦૮-૦૪-૨૦	૦૯-૦૪-૨૦	૧૦-૦૪-૨૦	૧૧-૦૪-૨૦	૧૨-૦૪-૨૦

વરસાદ(મી.મી.)	0	0	0	0	0
મહત્તમતાપમાન(સે.ગ્રે)	39	39	39	40	41
લઘુતમતાપમાન(સે.ગ્રે)	22	22	22	22	23
વાદળનીસ્થિતિ(ઓકટા)	0	0	0	0	1
મહત્તમભેજ (%)સવાર	80	78	85	76	55
લઘુતમભેજ (%) બપોરબાદ	25	34	28	26	23
પવનનીગતિ(કિમી/કલાક)	14	13	11	9	12
પવનનીદિશા(ડીગ્રી)	255	266	280	308	320

જામનગર: જીલ્લામાં આગામી તા.૦૮-૦૪-૨૦૨૦ થી તા.૧૨-૦૪-૨૦૨૦ દરમ્યાન હવામાન સુકું, ગરમ અને ચોખ્ખું રહેશે. પવનની દિશા પશ્ચિમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૯ થી ૧૬ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૭-૩૯ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૩-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ પેરામીટર	દિવસ-૧ ૦૮-૦૪-૨૦	દિવસ-૨ ૦૯-૦૪-૨૦	દિવસ-૩ ૧૦-૦૪-૨૦	દિવસ-૪ ૧૧-૦૪-૨૦	દિવસ-૫ ૧૨-૦૪-૨૦
વરસાદ(મી.મી.)	0	0	0	0	0
મહત્તમતાપમાન(સે.ગ્રે)	38	37	37	38	39
લઘુતમતાપમાન(સે.ગ્રે)	23	23	23	24	25
વાદળનીસ્થિતિ(ઓકટા)	0	0	0	0	0
મહત્તમભેજ (%)સવાર	83	85	89	83	66
લઘુતમભેજ (%) બપોરબાદ	46	53	49	45	39
પવનનીગતિ(કિમી/કલાક)	13	14	12	9	16
પવનનીદિશા(ડીગ્રી)	256	269	288	315	320

દેવભૂમિ દ્વારકા : જીલ્લામાં આગામી તા.૦૮-૦૪-૨૦૧૯ થી તા.૧૨-૦૪-૨૦૨૦ દરમ્યાન હવામાન સુકું, ગરમ અને ચોખ્ખું રહેશે. પવનની દિશા પશ્ચિમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૯ થી ૧૬ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૭-૩૯ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૩-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ પેરામીટર	દિવસ-૧ ૦૮-૦૪-૨૦	દિવસ-૨ ૦૯-૦૪-૨૦	દિવસ-૩ ૧૦-૦૪-૨૦	દિવસ-૪ ૧૧-૦૪-૨૦	દિવસ-૫ ૧૨-૦૪-૨૦
વરસાદ(મી.મી.)	0	0	0	0	0
મહત્તમતાપમાન(સે.ગ્રે)	38	37	37	38	39
લઘુતમતાપમાન(સે.ગ્રે)	23	23	23	24	25
વાદળનીસ્થિતિ(ઓકટા)	0	0	0	0	0
મહત્તમભેજ (%)સવાર	83	85	89	83	66
લઘુતમભેજ (%) બપોરબાદ	46	53	49	45	39
પવનનીગતિ(કિમી/કલાક)	13	14	12	9	16
પવનનીદિશા(ડીગ્રી)	256	269	288	315	320

અમરેલી : જીલ્લામાં આગામી તા.૦૮-૦૪-૨૦૨૦ થી તા.૧૨-૦૪-૨૦૨૦ દરમ્યાન હવામાન સુકું, ગરમ અને ચોખ્ખું રહેશે. પવનની દિશા પશ્ચિમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૧૧ થી ૧૩ કીમી/કલાક રહેની શક્યતા છે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૪૦-૪૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૩-૨૫ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ પેરામીટર	દિવસ-૧ ૦૮-૦૪-૨૦	દિવસ-૨ ૦૯-૦૪-૨૦	દિવસ-૩ ૧૦-૦૪-૨૦	દિવસ-૪ ૧૧-૦૪-૨૦	દિવસ-૫ ૧૨-૦૪-૨૦
વરસાદ(મી.મી.)	0	0	0	0	0
મહત્તમતાપમાન(સે.ગ્રે)	40	40	40	41	42




લઘુતમતાપમાન(સે.ગ્રે)	23	23	23	24	25
વાદળનીસ્થિતિ(ઓકટા)	0	0	0	0	0
મહતમભેજ (%)સવાર	78	80	76	71	58
લઘુતમભેજ (%) બપોરબાદ	29	36	31	33	27
પવનનીગતિ(કિમી/કલાક)	13	13	12	11	12
પવનનીદિશા(ડીગ્રી)	281	285	308	336	290


સુરેન્દ્રનગર : જીલ્લામાં આગામી તા.૦૪-૦૪-૨૦૨૦ થી તા.૦૮-૦૪-૨૦૨૦ દરમિયાન હવામાન સુકું, ગરમ અને મોટાભાગે યોખ્ખું રહેશે. પવનની દિશા પશ્ચિમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ **૧૨ થી ૧૫ કીમી/કલાક** રહેવાની શક્યતા છે .આ સમય ગાળામાં મહતમ તાપમાન દિવસ દરમિયાન **૩૯-૪૧** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમિયાન **૨૨-૨૩** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ પેરામીટર	દિવસ-૧ ૦૮-૦૪-૨૦	દિવસ-૨ ૦૯-૦૪-૨૦	દિવસ-૩ ૧૦-૦૪-૨૦	દિવસ-૪ ૧૧-૦૪-૨૦	દિવસ-૫ ૧૨-૦૪-૨૦
વરસાદ(મી.મી.)	0	0	0	0	0
મહતમતાપમાન(સે.ગ્રે)	39	39	39	40	41
લઘુતમતાપમાન(સે.ગ્રે)	22	22	22	22	23
વાદળનીસ્થિતિ(ઓકટા)	1	0	0	0	1
મહતમભેજ (%)સવાર	72	69	76	68	53
લઘુતમભેજ (%) બપોરબાદ	22	29	27	27	24
પવનનીગતિ(કિમી/કલાક)	15	14	12	12	15
પવનનીદિશા(ડીગ્રી)	261	269	286	311	329

- **૦૮ થી ૧૨-એપ્રિલ-૨૦૨૦** દરમિયાન હવામાન સુકું, ગરમ અને યોખ્ખું રહેશે. પવનની દિશા વાયવ્ય અને પશ્ચિમની મોટાભાગે રહેવાની અને પવનની ઝડપ **૯ થી ૧૬ કીમી/કલાક** રહેવાની શક્યતા છે .આ સમયગાળામાં મહતમ તાપમાન દિવસ દરમિયાન **૩૭-૪૨** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમિયાન **૨૨-૨૫** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.
- કોરોના વાયરસને ફેલાતો અટકાવવા બીનજરૂર અવાર જવર ટાળવી, કોઈપણ સંજોગોમાં સમુહમાં ભેગા થવું નહિ, વારંવાર સાબુથી હાથ ધોવા અને પછી જ આંખ, કાન અને મોને અડકવું. મોં પર કપડું બાંધી રાખવું, છીંક અને ઉધરસ વખતે મો અને નાક આડે રૂમાલ અવશ્ય રાખવો અને કાપણી, ગ્રેશિંગ અને અન્ય ખેતી વિષયક કામગીરી દરમિયાન સામાજીક અંતર (બે વ્યક્તિ વચ્ચે ૪ થી ૫ ફૂટ) જાળવવું તેમજ ભોજન અને આરામના સમયે અલગ અલગ દુર બેસવું.
- આગામી તા. ૧૨ થી ૧૬ એપ્રિલ દરમિયાન હવામાન યોખ્ખું અને ગરમ રહેશે. તેમજ લુ પડવાની શક્યતા હોય બપોરના સમયે તાપમાં જવું નહિ.
- આ હવામાન પરિસ્થિતિમાં ઉનાળુ શાકભાજી, તલ, અને ઘાસચારાના પાકોને સવારે અથવા સાંજના સમયે જરૂરીયાત મુજબ હળવું પિયત આપવું.
- શિયાળું પાકોને સંગ્રહ કરતા પહેલા તેના દાણાને વ્યવસ્થિત સફાઈ કરી, દાણામાં ૮ ટકા કરતા ઓછો ભેજ રહે ત્યાં સુધી સૂર્ય તાપમાં સુકવી પછી સંગ્રહ કરવો. તેમજ જે ગોડાઉનમાં સંગ્રહ કરવાનો હોય તે ગોડાઉનમાંથી સંગ્રહેલ જુના દાણાઓ બહાર કાઢી વ્યવસ્થિત સફાઈ કરવી તેમજ દીવાલોમાં પડેલ તિરાડોઓનું રીપેરીંગ કરવું અને ગોડાઉનમાં ૦.૫ ટકા મેલાથીયોન દવાના દ્રાવણનો છંટકાવ કરવો. દવાના છંટકાવ પછી ૭ થી ૮ દિવસ ગોડાઉન હવાયુસ્ત રીતે બંધ રાખવું. તેમજ દાણા ભરવા માટેના કોથળા સુર્યતાપમાં સૂકવવા જેથી જીવાતોના ઈંડા અને કોશેટાનો નાશ થાય.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટરપાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
-----------	------------	---------------	--

ઉનાળુ મગફળી	ફૂલ અને સુયા 	-	• ફૂલ અને સુયા અવસ્થાએ મગફળીને હળવું પિયત આપવું.
ઉનાળુ તલ	વૃદ્ધિ 	સફેદ માખી અને પાનવાળનાર (માથા બાંધનાર) ઈયળ	ઉનાળુ તલમાં શ્રીપ્સ અને પાનવાળનાર(માથા બાંધનાર) ઈયળના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મિલિ અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ડાયફેન્થાયુરોન ૫૦ ટકા વે.પા. ૧૬ ગ્રામ દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ તેમજ તલના પાકને ફૂલ અવસ્થાએ હળવું પિયત આપવું. ત્યારબાદ જમીનની પ્રત મુજબ ૧૦-૧૨ દિવસના ગાળે પિયત આપવું. વધારે પ્રમાણમાં ટૂંકા અંતરે પિયત આપવા નહિ. જરૂર કરતા વધારે ભેજથી તલમાં સૂકારો અને કાળીયો રોગ આવવાની શક્યતા રહે છે.
ઉનાળુ મગ અને અડદ	વૃદ્ધિ	શ્રીપ્સ અને સફેદ માખી	ઉનાળુ મગ અને અડદમાં શ્રીપ્સ અને સફેદ માખીના નિયંત્રણ માટે પ્રોફેનોફોસ ૨૦ મી.લી. અને ડાયફેન્થાયુરોન ૫૦ ટકા વે.પા. ૧૬ ગ્રામ અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
ઉનાળુ રીંગણ, ભીંડા, તુરીયા, દૂધી, ગલકા, કારેલા, ગુવાર, કાકડી, તરબુચ	વૃદ્ધિ	શ્રીપ્સ અને સફેદ માખી	ઉનાળુ રીંગણ, ભીંડા, તુરીયા, દૂધી, ગલકા, કારેલા, ગુવાર, કાકડી, તરબુચ અને તરબુચમાં શ્રીપ્સ અને સફેદ માખીના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મિલિ અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ડાયફેન્થાયુરોન ૫૦ ટકા વે.પા. ૧૬ ગ્રામ અથવા ટ્રાયઝોફોસ ૪૦ ઈસી ૨૫ મિ.લી. દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.
ડુંગળી	વૃદ્ધિ વિકાસ	શ્રીપ્સ, પાનનો સૂકારો 	➤ શ્રીપ્સના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦મિલિ અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને પ્રોફેનોફોસ ૩૦ મિલી અથવા લેમડા સાયહેલોથ્રીન ૧૫ મિલી દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. અતિ ઉપદ્રવ હોય તો સ્પીનોસાઈડ ૪ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ ડુંગળીના પાકમાં જાંબલી ધાબાના રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા કાર્બેન્ડાઝીમ ૧૦ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના/૧૦ લીટર પાણીમાં મિશ્ર કરી ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા. ➤ નાઈટ્રોજન યુક્ત એમોનિયમ સલ્ફેટ ખાતરનો પુરતી ડોઝ ૩૦ કીલો ગ્રામ પ્રતિ હે. આપવો.
રીંગણી અને ટમેટી	ફૂલ/ફળ	શ્રીપ્સ, સફેદ માખી ડુંખ અને ફળ કોરી ખાનાર ઈયળ	➤ શ્રીપ્સ અને સફેદ માખીના નિયંત્રણ માટે લીબોળીનું તેલ ૫૦ મિલિ અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને ડાયફેન્થાયુરોન ૫૦ ટકા વે.પા. ૧૬ ગ્રામ અથવા ટ્રાયઝોફોસ ૪૦ ઈસી ૨૫ મિ.લી. દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ ફળ અને ડોકાની ઈયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને ક્લોરોનટ્રાનીલીપ્રોલ ૧૮.૫

		પાન કથીરી	એસ.સી (રીનાક્ષીપાયર) દવા ૩ મિલી અથવા ડીડીવીપી ૭૬ ઈસી મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો. ➤ જો પાનકથીરીનો ઉપદ્રવ હોય તો પ્રોપરગાઈટ ૫૭ ઈ.સી. ૧૫ મિલી પ્રતિ પંપ છંટકાવ કરવો.
મરચી	વૃધ્ધિ/ફુલ/ફળ	થ્રીપ્સ, સફેદ માખી	➤ પ્રોફેનોફોસ ૨૦ મીલી અને લીબોળીનું તેલ ૫૦ મિલી અથવા લીબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. થ્રીપ્સનો વધુ ઉપદ્રવ હોય તો સ્પીનોસાદ અથવા સ્પીનેટોરમ દવા ૫ મિલી. પ્રતિ પંપ છાંટવી.
આંબો	ફૂલ		➤ આંબાને હળવું પિયત આપવું.
પશુ/મરઘા 	તાપમાન વધારે રહેવાના કારણે ઢોરને દિવસના સમયે છાયડામાં રાખવા અને મિનરલ મિક્શર યુક્ત પીવાના સ્વચ્છ પાણીની વ્યવસ્થા કરાવી. દુધાળા પશુઓને ઉનાળામાં ઠંડક આપવા પાણીનો છંટકાવ કરવો. ખોરાકમાં લીલા ચારનું પ્રમાણ વધારવું. દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા બોરિક પવડરથી બરાબર સાફ કરવું. દુધાળા પશુઓના આઉને ઝીંક ઓક્સાઈડ અથવા બોરિક પવડરથી બરાબર સાફ કરવું. પશુઓમાં ગળસુંઢો અને UF\9LIF તાવના રોગના નિયંત્રણ માટે રસીકરણ કરાવવું. ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાઝ ૨ મી.લી. ૧ લીટર પાણીમાં નાખીને છાંટવી. વિટામીન એ તથા ડી પુરકો આપવા. કરમિયાની દવા આપવી.		

જુક્યુ/ટેકનિકલ ઓફિસર).સુખે/(એગ્રોમેટ/બુલેટીન / / ૨૦૨૦
તરઘડિયા)રાજકોટ,(તારીખ: ૦૭/૦૪/૨૦૨૦.

પ્રતિ,
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનંતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર
સુકી ખેતીસંશોધન કેન્દ્ર
જુનાગઢ કૃષિ નિવર્સીટી
તરઘડીયા



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94269 38235



Agromet Advisory Service bulletin No.3-for the Rajkot district
(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
(Period 08 to 12-April - 2020)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	0.0
Maximum temperature (°C)	37.0-40.0
Minimum temperature (°C)	19.0-22.0
Relative Humidity (%) I	61-74
Relative Humidity (%) II	20-25
Wind speed (Kmph)	3.9-6.1


Weather forecast until 08-30 hrs. of 12/04/2020





(1) RAJKOT DISTRICT:

	08/04/20	09/04/20	10/04/20	11/04/20	12/04/20
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	39	39	39	40	41
Min.Temp.(°C)	22	22	22	22	23
Total CC (octa)	0	0	0	0	1
RH-I (%)	80	78	85	76	55
RH-II (%)	25	34	28	26	23
Wind Speed (kmph)	14	13	11	9	12
Wind Direction(deg.)	255	266	280	308	320

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none">• Rain fall: Dry, hot and clear weather forecasted next five days.• Temperature: Maximum and minimum temperature likely to be in range of 39-41°C and 22-23°C, respectively.• Relative Humidity: The RH-I and RH-II is likely to be in the range of 55-85% & 23-34, respectively.• Wind Direction and Speed: Direction and speed of wind is likely to be W, NW and 9-14 km/h, respectively.• Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with
-----------------------	--

	<p>soap at reasonable interval, social distancing during harvesting and other agricultural operation in all crops.</p> <ul style="list-style-type: none"> Weather outlook for subsequent week: Dry, clear and hot weather very likely to observe during 13-16 April-2020. Due to prevailing hot weather irrigate summer vegetable and fodder crops during night hours. Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop. Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed. 		
Crops	Stage	Insect/Pest/ Variety	Advisory
Summer Groundnut	Flowering/ pegging		Apply light irrigation at flowering stage
Summer Sesame	Flowering initiation	White fly/ Leaf Webber	For the control thrips and leaf webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Green gram & Black gram	Growing	White fly & Thrips	For the control thrips in green & black gram spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water.
Brinjal, Lady finger, Ridge gourd, Bottle gourd, Sponge guard, Bittergourd, Cluster bean, Water melon	Growing	Thrips & white fly	Monitor the crops for infestation of thrips & white fly in brinjal, lady finger, ridge, sponge, bottle and bitter gourd, cluster bean and water melon, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % WP @ 16 gm /10 lit of water.

Onion	Growth/harvesting	Thrips Leaf blight→ 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Make alternate spray of Mancozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.
Brinjal & Tomato	Flowering/fruiting	Sucking pest → Shoot & Fruit borer  Mites	Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer. If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.
Chili	Flowering & fruiting	Thrips → 	Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.
Mango	Fruit development	-	To prevent dropping of immature fruits in mango spray 2 gm NAA and 2 kg Urea in 100 litres of water. Apply irrigation as per need.
Livestock & poultry 	Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day .Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.		

Technical Officer, GKMS Unit, DFRS, JAU, Targadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-3 for the Morbi district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 08 to 12-April - 2020)

Weather forecast until 08-30 hrs. Of 12/04/2020



(2) MORBI DISTRICT:




	08/04/20	09/04/20	10/04/20	11/04/20	12/04/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	39	39	39	40	41
Min. Temp.(°C)	22	22	22	22	23
Total CC (octa)	0	0	0	0	1
RH-I (%)	80	78	85	76	55
RH-II (%)	25	34	28	26	23
Wind Speed (kmph)	14	13	11	9	12
Wind Direction(deg.)	255	266	280	308	320

Agro meteorological Advisories:-

General Advice

- **Rain fall: Dry, hot and clear weather forecasted next five days.**
- **Temperature:** Maximum and minimum temperature likely to be in range of 39-41°C and 22-23°C, respectively.
- **Relative Humidity:** The RH-I and RH-II is likely to be in the range of 55-85% & 23-34, respectively.
- **Wind Direction and Speed:** Direction and speed of wind is likely to be W, NW and 9-14 km/h, respectively.
- **Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing during harvesting and other agricultural operation in all crops.**
- **Weather outlook for subsequent week: Dry, clear and hot weather very likely to observe during 13-16 April-2020.**
- **Due to prevailing hot weather irrigate summer vegetable and fodder crops during night hours.**
- Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop.

	Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed.		
Crops	Stage	Insect/Pest/ Variety	Advisory
Summer Groundnut	Flowering/ pegging		Apply light irrigation at flowering stage
Summer Sesame	Flowering initiation	White fly/ Leaf Webber	For the control thrips and leaf webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Green gram & Black gram	Growing	White fly & Thrips	For the control thrips in green & black gram spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water
Brinjal, Lady finger, Ridge gourd, Bottle gourd, Sponge guard, Bittergourd, Cluster bean, Water melon	Growing	Thrips & white fly	Monitor the crops for infestation of thrips & white fly in brinjal, lady finger, ridge, sponge, bottle and bitter gourd, cluster bean and water melon, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % WP @ 16 gm /10 lit of water.
Onion	Growth/harve sting	Thrips Leaf blight→ 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Make alternate spray of Mancozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.
Brinjal& Tomato	Flowering/ fruiting	Sucking pest → Shoot & Fruit borer	Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of

		 <p>Mites</p>	<p>chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</p>
Chili	Flowering & fruiting	<p>Thrips →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p>
Mango	Fruit development	-	<p>To prevent dropping of immature fruits in mango spray 2 gm NAA and 2 kg Urea in 100 litres of water. Apply irrigation as per need.</p>
Livestock & poultry 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day .Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-3 for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 08 to 12-April- 2020)

Weather forecast until 08-30 hrs. Of 12/04/2020



(3) JAMNAGAR DISTRICT:




	08/04/20	09/04/20	10/04/20	11/04/20	12/04/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	38	37	37	38	39
Min. Temp.(°C)	23	23	23	24	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	83	85	89	83	66
RH-II (%)	46	53	49	45	39
Wind Speed (kmph)	13	14	12	9	16
Wind Direction(deg.)	256	269	288	315	320

Agro meteorological Advisories:-

General Advice

- **Rain fall: Dry, hot and clear weather forecasted next five days.**
- **Temperature:** Maximum and minimum temperature likely to be in range of 37-39°C and 23-25°C, respectively.
- **Relative Humidity:** The RH-I and RH-II is likely to be in the range of 66-89% & 39-53, respectively.
- **Wind Direction and Speed:** Direction and speed of wind is likely to be W, NW and 9-16 km/h, respectively.
- **Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing during harvesting and other agricultural operation in all crops.**
- **Weather outlook for subsequent week: Dry, clear and hot weather very likely to observe during 13-16 April-2020.**
- **Due to prevailing hot weather irrigate summer vegetable and fodder crops during night hours.**
- Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop.

	Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed.		
Crops	Stage	Insect/Pest/ Variety	Advisory
Summer Groundnut	Flowering/ pegging		Apply light irrigation at flowering stage
Summer Sesame	Flowering initiation	White fly/ Leaf Webber	For the control thrips and leaf webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Green gram & Black gram	Growing	White fly & Thrips	For the control thrips in green & black gram spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water.
Brinjal, Lady finger, Ridge gourd, Bottle gourd, Sponge guard, Bittergourd, Cluster bean, Water melon	Growing	Thrips & white fly	Monitor the crops for infestation of thrips & white fly in brinjal, lady finger, ridge, sponge, bottle and bitter gourd, cluster bean and water melon, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % WP @ 16 gm /10 lit of water.
Onion	Growth/harve sting	Thrips Leaf blight→ 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Make alternate spray of Mancozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.
Brinjal& Tomato	Flowering/ fruiting	Sucking pest → Shoot &Fruit borer	Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of

		 <p>Mites</p>	<p>chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</p>
Chili	Flowering & fruiting	<p>Thrips →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p>
Mango	Fruit development	-	<p>To prevent dropping of immature fruits in mango spray 2 gm NAA and 2 kg Urea in 100 litres of water.</p> <p>Apply irrigation as per need.</p>
Livestock & poultry 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day .Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-3 for the Devbhumi Dwarka district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 08 to 12-April- 2020)

Weather forecast until 08-30 hrs. of 12/04/2020



(4) DEVBHUMI DWARKA DISTRICT:




	08/04/20	09/04/20	10/04/20	11/04/20	12/04/20
Rainfall (mm)	0	0	0	0	0
Max. Temp.(°C)	38	37	37	38	39
Min. Temp.(°C)	23	23	23	24	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	83	85	89	83	66
RH-II (%)	46	53	49	45	39
Wind Speed (kmph)	13	14	12	9	16
Wind Direction(deg.)	256	269	288	315	320

Agro meteorological Advisories:-

General Advice

- **Rain fall: Dry, hot and clear weather forecasted next five days.**
- **Temperature:** Maximum and minimum temperature likely to be in range of 37-39°C and 23-25°C, respectively.
- **Relative Humidity:** The RH-I and RH-II is likely to be in the range of 66-89% & 39-53, respectively.
- **Wind Direction and Speed:** Direction and speed of wind is likely to be W, NW and 9-16 km/h, respectively.
- **Keeping in view of Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing during harvesting and other agricultural operation in all crops.**
- **Weather outlook for subsequent week: Dry, clear and hot weather very likely to observe during 13-16 April-2020.**
- **Due to prevailing hot weather irrigate summer vegetable and fodder crops during night hours.**
- Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop.

	Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed.		
Crops	Stage	Insect/Pest/ Variety	Advisory
Summer Groundnut	Flowering/ pegging		Apply light irrigation at flowering stage
Summer Sesame	Flowering initiation	White fly/ Leaf Webber	For the control thrips and leaf webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Green gram & Black gram	Growing	White fly & Thrips	For the control thrips in green & black gram spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water.
Brinjal, Lady finger, Ridge gourd, Bottle gourd, Sponge guard, Bittergourd, Cluster bean, Water melon	Growing	Thrips & white fly	Monitor the crops for infestation of thrips & white fly in brinjal, lady finger, ridge, sponge, bottle and bitter gourd, cluster bean and water melon, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % WP @ 16 gm /10 lit of water.
Onion	Growth/harve sting	Thrips Leaf blight→ 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Make alternate spray of Mancozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.
Brinjal& Tomato	Flowering/ fruiting	Sucking pest → Shoot & Fruit borer	Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of

		 <p>Mites</p>	<p>chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</p>
Chili	Flowering & fruiting	<p>Thrips →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p>
Mango	Fruit development	-	<p>To prevent dropping of immature fruits in mango spray 2 gm NAA and 2 kg Urea in 100 litres of water.</p> <p>Apply irrigation as per need.</p>
Livestock & poultry 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day .Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :9426938235



Agro met Advisory Service bulletin No.-3 for the Amreli district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 08 to 12-April - 2020)

Weather forecast until 08-30 hrs. of 12/04/2020



(5) AMRELI DISTRICT:




	08/04/20	09/04/20	10/04/20	11/04/20	12/04/20
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	40	40	40	41	42
Min.Temp.(°C)	23	23	23	24	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	78	80	76	71	58
RH-II (%)	29	36	31	33	27
Wind Speed (kmph)	13	13	12	11	12
Wind Direction(deg.)	281	285	308	336	290

Agro meteorological Advisories:

General Advice

- **Rain fall: Dry, hot and clear weather forecasted next five days.**
- **Temperature:** Maximum and minimum temperature likely to be in range of 40-42°C and 23-25°C, respectively.
- **Relative Humidity:** The RH-I and RH-II is likely to be in the range of 58-80% & 27-36, respectively.
- **Wind Direction and Speed:** Direction and speed of wind is likely to be W, NW and 11-13 km/h, respectively.
- **Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing during harvesting and other agricultural operation in all crops.**
- **Weather outlook for subsequent week: Dry, clear and hot weather very likely to observe during 13-16 April-2020.**
- **Due to prevailing hot weather irrigate summer vegetable and fodder crops during night hours.**
- Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop.

	Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed.		
Crops	Stage	Insect/Pest/ Variety	Advisory
Summer Groundnut	Flowering/ pegging		Apply light irrigation at flowering stage
Summer Sesame	Flowering initiation	White fly/ Leaf Webber	For the control thrips and leaf webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Green gram & Black gram	Growing	White fly & Thrips	For the control thrips in green & black gram spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water.
Brinjal, Lady finger, Ridge gourd, Bottle gourd, Sponge guard, Bittergourd, Cluster bean, Water melon	Growing	Thrips & white fly	Monitor the crops for infestation of thrips & white fly in brinjal, lady finger, ridge, sponge, bottle and bitter gourd, cluster bean and water melon, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % WP @ 16 gm /10 lit of water.
Onion	Growth/harve sting	Thrips Leaf blight→ 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Make alternate spray of Mancozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.
Brinjal& Tomato	Flowering/ fruiting	Sucking pest → Shoot &Fruit borer	Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of

		 <p>Mites</p>	<p>chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</p>
Chili	Flowering & fruiting	<p>Thrips →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p>
Mango	Fruit development	-	<p>To prevent dropping of immature fruits in mango spray 2 gm NAA and 2 kg Urea in 100 litres of water.</p> <p>Apply irrigation as per need.</p>
Livestock & poultry 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day .Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell:9426938235



Agro met Advisory Service bulletin No.-3 for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 08 to 12-April- 2020)

Weather forecast until 08-30 hrs of 12/04/2020



(6) SURENDRANAGAR DISTRICT:




	08/04/20	09/04/20	10/04/20	11/04/20	12/04/20
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	39	39	39	40	41
Min.Temp.(°C)	22	22	22	22	23
Total CC (octa)	1	0	0	0	1
RH-I (%)	72	69	76	68	53
RH-II (%)	22	29	27	27	24
Wind Speed (kmph)	15	14	12	12	15
Wind Direction(deg.)	261	269	286	311	329

Agro meteorological advisories:

General Advice

- **Rain fall: Dry, hot and Mainly clear to clear weather forecasted next five days.**
- **Temperature:** Maximum and minimum temperature likely to be in range of 39-41°C and 22-23°C, respectively.
- **Relative Humidity:** The RH-I and RH-II is likely to be in the range of 53-76% & 22-29, respectively.
- **Wind Direction and Speed:** Direction and speed of wind is likely to be W, NW and 12-15 km/h, respectively.
- **Keeping in view of threat of COVID-19 spread farmers are advised to follow the government guidelines, personal hygiene, use mask, ensure hand washing with soap at reasonable interval, social distancing during harvesting and other agricultural operation in all crops.**
- **Weather outlook for subsequent week: Dry, clear and hot weather very likely to observe during 13-16 April-2020.**
- **Due to prevailing hot weather irrigate summer vegetable and fodder crops during night hours.**
- Before storage of rabi crops produce, grains should be properly cleaned and it should be dried properly to the recommended level of moisture content. Godowns should be properly cleaned by removing all the grains and other storage matter of previous crop.

	Cracks and crevices of the walls should be cleaned and repaired and whitewashing should be done. Those who can afford are advised to spray 0.5 % malathion solution and keep the chamber closed for 7-8 days. The gunny bags should be dried in the sun so that eggs and pupating stages of insects as well as inoculum of diseases get destroyed.		
Crops	Stage	Insect/Pest/ Variety	Advisory
Summer Groundnut	Flowering/ pegging		Apply light irrigation at flowering stage
Summer Sesame	Flowering initiation	White fly/ Leaf Webber	For the control thrips and leaf webber spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water. To prevent Phytophthora blight and wilt do not over irrigate sesame, apply light irrigation as per need only.
Green gram & Black gram	Growing	White fly & Thrips	For the control thrips in green & black gram spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with difenthiuron 16 gm/10 lit of water.
Brinjal, Lady finger, Ridge gourd, Bottle gourd, Sponge guard, Bittergourd, Cluster bean, Water melon	Growing	Thrips & white fly	Monitor the crops for infestation of thrips & white fly in brinjal, lady finger, ridge, sponge, bottle and bitter gourd, cluster bean and water melon, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % WP @ 16 gm /10 lit of water.
Onion	Growth/harve sting	Thrips Leaf blight→ 	For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval. Make alternate spray of Mancozeb 25 gm and Carbendazim 10gm in 10 liters of water at 10 days interval.
Brinjal& Tomato	Flowering/ fruiting	Sucking pest → Shoot & Fruit borer	Monitor the crop for infestation of white fly in brinjal, if require spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with diafenthiuron 50 % wp @ 16 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of

		 <p>Mites</p>	<p>chlorantraniliprole 18.5 % SC 3.0 ml or DDVP 7 ml/10 litre of water is advised against fruit borer.</p> <p>If infestation is found, then make spraying of Propargite 57 % EC @ 10 ml/ 10 litre of water for the control of Mites under clear sky.</p>
Chili	Flowering & fruiting	<p>Thrips →</p> 	<p>Under present weather conditions severe attack of thrips has been reported. Spray profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of heavy attack make spray of spinosad or spinetoram 3 ml per 10 liter of water.</p>
Mango	Fruit development	-	<p>To prevent dropping of immature fruits in mango spray 2 gm NAA and 2 kg Urea in 100 litres of water.</p> <p>Apply irrigation as per need.</p>
Livestock & poultry 	<p>Under present weather condition keep animals under shade or in sheds during noon hours, provide plenty of cool water mixed with minerals for drinking and shower the animals with cold water two to three times in a day .Do Vaccination for Haemorrhagic septicemia (H.S.) disease and Blue Quarter (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.</p>		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.