



ગ્રામીણ કૃષિ મૌસમ સેવા  
હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન  
ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ  
તા.14-01-2017 થી તા.18-01-2017



(ગ્રામીણ કૃષિ મૌસમસેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કૃ.યુ., તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારાસંપન્ન.)

કૃષિહવામાન બુલેટીન નં. **83/2016**

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

હવામાન પરિબલો	
વરસાદ(મી.મી)	0.0
મહત્તમ તાપમાન (સે.ગ્રે)	24.3-26.1
લઘુત્તમ તાપમાન (સે.ગ્રે)	7.6-10.0
ભેજ(%) સવાર	45-75
ભેજ(%) બપોરબાદ	45-56
પવનની ગતિ (કિમી/કલાક)	4.1-5.7

હવામાન આગાહી: પત્રક-ક

ઉત્તર સૌરાષ્ટ્રખેત આબોહવાકીય વિભાગહવામાન આગાહી

તા.14-01-2017 થી 18-01-2017

કૃષિ સલાહ:

સામાન્ય	<p>➤ ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના રાજકોટ</p> <p><b>રાજકોટ:</b> જલ્દામાં આગામી તા.14-01-2017 થી તા.18-01-2017 દરમિયાન હવામાન સુકું અને અંશતઃ વાદળછાયુ રહેશે. પવનની દિશા ઈશાન, નૈઋત્ય અને ઉત્તરની રહેવાની અને પવનની ઝડપ <b>15 થી 21 કીમી/કલાક</b> રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન <b>27-28</b> ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન <b>12-14</b> ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.</p>					
	તારીખ	દિવસ-1	દિવસ-2	દિવસ-3	દિવસ-4	દિવસ-5
	પેરામીટર	14-01-17	15-01-17	16-01-17	17-01-17	18-01-17
	વરસાદ(મી.મી)	0	0	0	0	0
	મહત્તમ તાપમાન (સે.ગ્રે)	27	27	27	28	28
	લઘુત્તમ તાપમાન (સે.ગ્રે)	12	13	14	12	12

વાદળનીસ્થિતિ(ઓકટા)	0	2	3	4	0
મહતમ ભેજ (%) સવાર	28	43	45	51	49
લઘુતમ ભેજ (%) બપોરબાદ	17	21	25	31	24
પવનની ગતિ મી/કલાક	15	12	19	20	21
પવનની દિશા (ડીગ્રી)	67	79	225	19	22

**જામનગર:** જીલ્લામાં આગામી તા .14-01-2017 થી તા .18-01-2017 દરમ્યાન હવામાન સુકું અને અંશતઃ વાદળછાયુ રહેશે. પવનની દિશા પૂર્વ, અગ્નિ અને ઈશાનની રહેવાની અને પવનની ઝડપ **15 થી 23 કીમી/કલાક** રહેવાની શક્યતા છે. આ સમયગાળામાં મહતમ તાપમાન દિવસ દરમ્યાન **27-28** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન **12-14** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-1	દિવસ-2	દિવસ-3	દિવસ-4	દિવસ-5
પેરામીટર	14-01-17	15-01-17	16-01-17	17-01-17	18-01-17
વરસાદ(મી.મી)	0	0	0	0	0
મહતમ તાપમાન (સે.ગ્રે)	27	27	27	28	28
લઘુતમ તાપમાન (સે.ગ્રે)	12	13	14	12	12
વાદળની સ્થિતિ(ઓકટા)	0	3	4	4	0
મહતમ ભેજ (%) સવાર	31	44	47	53	52
લઘુતમ ભેજ (%) બપોરબાદ	19	22	29	34	28
પવનની ગતિ (કિમી/કલાક)	15	16	23	21	23
પવનની દિશા (ડીગ્રી)	75	36	123	26	28

**અમરેલી:** જીલ્લામાં આગામી તા. 14-01-2017 થી તા .18-01-2017 દરમ્યાન હવામાન સુકું અને અંશતઃ વાદળછાયુ રહેશે. પવનની દિશા ઈશાન, અગ્નિ, પશ્ચિમ અને પૂર્વની રહેવાની અને પવનનીઝડપ **9 થી 20 કીમી/કલાક** રહેવાની શક્યતા છે. આ સમયગાળામાં મહતમ તાપમાન દિવસ દરમ્યાન **26-28** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન **8-11** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-1	દિવસ-2	દિવસ-3	દિવસ-4	દિવસ-5
પેરામીટર	14-01-17	15-01-17	16-01-17	17-01-17	18-01-17
વરસાદ(મી.મી)	0	0	0	0	0
મહતમ તાપમાન (સે.ગ્રે)	26	26	27	27	28
લઘુતમ તાપમાન (સે.ગ્રે)	9	11	11	9	8
વાદળની સ્થિતિ(ઓકટા)	0	1	3	4	1
મહતમ ભેજ (%) સવાર	32	45	48	48	45


લઘુતમ ભેજ (%) બપોરબાદ	22	27	28	31	25
પવનની ગતિ કિમી/કલાક	16	9	17	20	20
પવનની દિશા (ડીગ્રી)	56	139	261	76	76




**સુરેન્દ્રનગર:** જીલ્લામાં આગામી તા .14-01-2017 થી તા .18-01-2017 દરમ્યાન હવામાન સુકું અને અંશતઃ વાદળછાયુ રહેશે. પવનની દિશા પૂર્વ, વાયવ્ય અને ઈશાનની રહેવાની અને પવનની ઝડપ **11 થી 18 કીમી/કલાક** રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન **27-28** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન **12-14** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.



તારીખ પેરામીટર	દિવસ-1 14-01-17	દિવસ-2 15-01-17	દિવસ-3 16-01-17	દિવસ-4 17-01-17	દિવસ-5 18-01-17
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમ તાપમાન (સે.ગ્રે)	27	27	27	28	28
લઘુતમ તાપમાન (સે.ગ્રે)	12	13	14	12	12
વાદળની સ્થિતિ(ઓકટા)	0	2	3	5	0
મહત્તમ ભેજ (%) સવાર	27	41	44	50	48
લઘુતમ ભેજ (%) બપોરબાદ	15	18	22	26	23
પવનની ગતિ કિમી/કલાક	16	11	18	16	17
પવનની દિશા (ડીગ્રી)	80	76	319	42	50






**18 જાન્યુઆરિ સુધી** હવામાન સુકું અને અંશતઃ વાદળછાયુ રહેશે. પવનની દિશા ઈશાન, ઉત્તર અને પુર્વની રહેવાની અને પવનની ઝડપ **11 થી 24 કીમી/કલાક** રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન **26-28** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન **9-13** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

- અમરેલિ જીલ્લામાં ઠંડી વધવાની શક્યતા હોય નીચા તાપમાનને કારણે શાકભાજી અને અન્ય પાકોને હિમની અસર ન થાય તે માટે સાંજના સમયે પિયત આપવું, પાકના આવશેસોનું આવરણ કરવું તેમજ ખેતર ફરતે રાત્રીના સમયે ધુમાડો કરવો.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
લસણ	ગાંજો બન્યાવો	શ્રીપ્સ, પાનની ટોચનો 	શ્રીપ્સના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને પ્રોફેનોફોસ ૨૦ મિલી અથવા લેમ્બડા સાયહેલોથ્રીન ૧૫ મિલી દવાને ૧૦

		<p>સુકારો</p> 	<p>લીટર પાણીમાં મિશ્ર લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</p> <p>પાનની ટોચનો સુકારાના નિયંત્રન માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામ ૧૦ લીટર પાણીમાં કરીને ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.</p>
ચણા	કુલ/પોપટા બન્ધાવા/ દાણા ભરાવા	લીલી અને લશ્કરી ઈયળ	<p>ચણાના પાકમાં લીલી અને લશ્કરી ઈયળના નિયંત્રણ માટે પક્ષીઓને બેસવા માટે 'T' આકારના બેલાખડા મુકવા. જો વધુ ઉપદ્રવ જણાય તો એમામેક્ટીન બેન્ઝોએટ ૬.૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.</p>
ધઉં	કુલ/દાણા ભરાવા	-	સમયસર પિયત આપવું.
જીરું	કુલ અને દાણા બન્ધાવા	<p>શ્રીપ્સ અને મોલો</p> <p>ચરમી અને રાખોડી</p>  <p>સુકારો</p>	<p>શ્રીપ્સ અને મોલોના ના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.</p> <p>તેમજ ચરમીનો ઉપદ્રવ ન થાય તે માટે ડાયથેન એમ ૪૫, ૨૭ ગ્રામ અને હેક્ઝાકોનાઝોલ ૧૫ મિલી દવા વારાફરતી ૧૦ લીટર પાણીમાં મિશ્ર કરીને ૧૫ દિવસના અંતરે છંટકાવ કરવો.</p>  <p>ઉભા પાકમાં સુકારાના નિયંત્રણ માટે પિયત આપવાનું બંધ કરવું.</p>

			
ધાણા	કુલ અને દાણા બન્ધાવા	થ્રીપ્સ અને મોલો	થ્રીપ્સ અને મોલોના ના નિયંત્રણ માટે લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને થાયોમીથોક્ઝામ ૫ ગ્રામ ૧૦ લીટર પાણીમાં કરીને છંટકાવ કરવો.
ડુંગળી	વૃદ્ધિ	જાંબલી ધાબા 	ડુંગળીના પાકમાં જાંબલી ધાબાના રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામ ૧૦ લીટર પાણીમાં કરીને ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.
રીંગણી અને ટમેટી	કુલ/ફળ	સફેદ માખી  ડુંખ અને ફળ કોરી ખાનાર ઈયળ →  કથીરી →	<ul style="list-style-type: none"> <li>લીંબોળીનું તેલ ૫૦મિલિ અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</li> <li>ફળ અને ડોકાની ઈયળોનો ઉપદ્રવ હોય તો નુકશાન પામેલ ફળ અને ડોકા તોડીને જમીનમાં દાંટી દેવા અને રીનાક્ષીપાયર દવા ૩ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</li> <li>રીંગણી અને ટમેટી ના પાકમાં પાન કથીરીના નિયંત્રણ માટે પ્રોપરગાઈટ દવા ૧૦ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરીને છંટકાવ કરવો.</li> </ul>
કોબી અને	વૃદ્ધિ/દડા	હિરાફૂદુ	કોબી અને ફલાવરમાં હિરાફૂદાણા નિયંત્રણ માટે સ્પીનોસાઈ ૩ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરી

ફલાવર			છંટકાવ કરવો
મરચી	ફુલ/ફળ	<p>પાનની કુકડ-થ્રીપ્સ→</p>  <p>કાલવ્રણ →</p> 	<p>પ્રોફેનોફોસ ૨૦ મી.લી.અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> <p>કાલવ્રણ રોગના નિયંત્રણ માટે મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.</p>
આંબો	ફૂલ	<p>મધીયો અને મગીયાની ઈયળ</p> <p>ભુકીછારો/રાખોડી</p>	<p>આંબામાં મધીયો અને મગીયાની ઈયળના નિયંત્રણ માટે ઇમીડાક્લોપ્રીડ ૫ મિલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p>  <p>આંબામાં ભુકીછારો/રાખોડીના નિયંત્રણ માટે હેક્ઝાકોનાઝોલ ૧૫ મિલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.</p> 
<p>5X]qDZ 3F</p> <p>પશુ/મર ઘા.</p> 	<p>નાના વાછરડાઓને રાત્રે ઠંડીથી બચવા પશુ ઘરમાં રખવા. પ્રજીવથી થતા બૃસેલ્લા અને અન્ય રોગોની ચકાસણી કરાવવી. ખરવા મોવાસાના રોગનું રસીકરણ કરાવવું. ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાઝ ૨ મી.લી. ૧ લીટર પાણીમાં નાખીને છાંટવી. વિટામીન એ તથા ડી પુરકો આપવા. કરમિયાની દવા આપવી તથા ઉછરતા પશુઓમાં બૃસેલ્લા કાફવુંડનું રસીકરણ કરાવવું. મરઘા ઘરના પશ્ચિમ દિશામાં</p>		

કંતાનના પડદા લગાવવા.

જુકૃયુ/ટેકનિકલ ઓફિસર.(સુખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૧૬, તરઘડિયા(રાજકોટ), તારીખ: 11/01/2017.

પ્રતિ,  
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,  
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમ માં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર  
સુકી ખેતી સંશોધન કેન્દ્ર  
જુનાગઢ કૃષિ યુનિવર્સિટી  
તરઘડીયા





**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :94277 28523**



**Agro met Advisory Service bulletin No.-83**  
**for the Rajkot district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)  
 (Period 14-18 January- 2017)

**Significant past weather for the preceding week of Rajkot**

Rainfall (mm)	0.0
Maximum temperature (°C )	24.3-26.1
Minimum temperature (°C )	7.6-10.0
Relative Humidity (%) I	45-75
Relative Humidity (%) II	45-56
Wind speed (Kmph)	4.1-5.7

**Weather forecast until 08-30 hrs of 18/01/2017**







**(1) RAJKOT DISTRICT:**





	14/01/17	15/01/17	16/01/17	17/01/17	18/01/17
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	27	27	27	28	28
Min.Temp.(°C)	12	13	14	12	12
Total CC (octa)	0	2	3	4	0
RH-I (%)	28	43	45	51	49
RH-II (%)	17	21	25	31	24
Wind Speed (kmph)	15	12	19	20	21
Wind Direction(deg.)	67	79	225	19	22



**Agro meteorological Advisories:**

<b>General Advice</b>	<ul style="list-style-type: none"> <li>• <b>Rain fall:</b> Dry and clear to partly cloudy weather forecasted for next five days.</li> <li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 27-28°C and 12-13°C, respectively.</li> <li>• <b>Relative Humidity:</b> The RH is likely to be in the range of 17-51 %</li> <li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be NE, E, SW, N and 15-21 km/h, respectively.</li> </ul>		
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/Variety</b>	<b>Advisory</b>
<b>Garlic</b>	Girthing	Thrips and	<b>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</b>



		 Alternaria leaf blight 	The disease infestation observed in standing crop. Increase periods between irrigations means do not over irrigate the crop. Make two Spray 27 gm Mencozeb/10 liters of water at 10 days interval.
<b>Chick pea</b>	Flowering/ pod formation	Monitor the crop regularly for and heliothes infestation	“T” shaped bird perches in and around crop field to be installed for controlling insect population. If infestation cross ETL spray 3 ml <b>SPINOSAD</b> in 10 litres of water.
<b>Wheat</b>	Milking	-	Apply irrigation without stress.
<b>Cumin</b>	Flowering /grain formation	<b>Aphid and thrips</b>  <b>Blight and Powery mildew</b>   <b>Wilt</b> 	Under present weather condition attack of thrips and aphids has been reported spray <b>imidacloprid 5 ml</b> /10 liters of water.  Under present weather condition spray <b>Hexaconazole</b> 15 ml/10 liters of water to prevent blight and powdery mildew infestation.   Stop irrigation to prevent further infection in standing crop. 
<b>Coriander</b>	Flowering & grain formation	Aphid, Thrips and Jassids	Spray <b>neem seed extract</b> 5% and <b>Thiomethoxam 5 gm/ 10 liters of water.</b> Under present weather condition spray <b>Hexaconazole</b> 15 ml/10 liters of water to prevent powdery mildew infestation.
<b>Onion</b>	Growth	Thrips	<b>For thrips control spray neem seed extract 5%</b>

		Purple blotch → 	and <b>Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</b>  Spray <b>Mencozeb</b> 25 gm and <b>hexaconazole</b> 15 ml in 10 liters of water at 10 days interval.
<b>Brinjal &amp; Tomato</b>	Flowering/ fruiting	<b>Sucking pest</b> →  & <b>Fruit borer</b>	To control sucking pest spray <b>neem seed extract</b> 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
<b>Cabbage and cauliflower</b>	Growth & fruiting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.
<b>Chili</b>	Flowering & fruiting	Thrips, →  Wilting → Anthracnose → 	In present weather conditions severe attack of thrips has been reported. Spray 3 ml <b>spinosad</b> in 10 litres of water.  Drenching of Trichoderma viridi 4 gm in 1 liter of water.  Spray <b>Mencozeb</b> 27 gm in 10 liters of water
<b>Mango</b>	Flowering	<b>Blossom midge and Leaf hopper</b> →  <b>Powdery</b>	To prevent damage spray <b>imidacloprid</b> 5 ml/10 litres of water.  Spray 15 ml <b>Hexaconzole</b> in 10 liters of water.

		<b>Mildew</b> → 	
<b>Livestock &amp; poultry</b> 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose. Put curtains on west side of poultry house.		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :94277 28523**



**Agro met Advisory Service bulletin No.-83 for the Jamnagar district**



(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)  
 (Period 14-18-Jan.- 2017)






**Weather forecast until 08-30 hrs of 18/01/2017**






**(2) JAMNAGAR DISTRICT:**

	14/01/17	15/01/17	16/01/17	17/01/17	18/01/17
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	27	27	27	28	28
Min.Temp.(°C)	12	13	14	12	12
Total CC (octa)	0	3	4	4	0
RH-I (%)	31	44	47	53	52
RH-II (%)	19	22	29	34	28
Wind Speed (kmph)	15	16	23	21	23
Wind Direction(deg.)	75	36	123	26	28

**Agro meteorological Advisories:-**

<b>General Advice</b>	<ul style="list-style-type: none"> <li>• <b>Rain fall:</b> Dry and clear to partly cloudy weather forecasted for next five days.</li> <li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 27-28°C and 12-14°C, respectively.</li> <li>• <b>Relative Humidity:</b> The RH is likely to be in the range of 19-53%</li> <li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be E, NE, SE, NE and 15-23 km/h, respectively.</li> </ul>		
Crops	Stage	Insect/Pest/Variety	Advisory
<b>Garlic</b>	Girthing	Thrips and  Alternaria leaf blight 	<p><b>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</b></p> <p>The disease infestation observed in standing crop. Increase periods between irrigations means do not over irrigate the crop.            Make two Spray 27 gm Mencozeb/10 liters of water at 10 days interval.</p>
<b>Chick pea</b>	Flowering/pod	Monitor the crop regularly for	“T” shaped bird perches in and around crop field to be installed for controlling insect population.

	formation	prodenia heliothis infestation and	If infestation cross ETL spray 3 ml <b>SPINOSAD</b> in 10 litres of water.
<b>Wheat</b>	Milking	-	Apply irrigation without stress.
<b>Cumin</b>	Flowering /grain formation	<b>Aphid and thrips</b>  <b>Blight and Powery mildew</b>    <b>Wilt</b>  	<p>Under present weather condition attack of thrips and aphids has been reported spray <b>imidacloprid 5 ml</b>/10 liters of water.</p> <p>Under present weather condition spray <b>Hexaconazole</b> 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
<b>Coriander</b>	Flowering & grain formation	Aphid, Thrips and Jassids	<b>Spray neem seed extract 5% and Thiomethoxam 5 gm/ 10 liters of water.</b> Under present weather condition spray <b>Hexaconazole</b> 15 ml/10 liters of water to prevent powdery mildew infestation.
<b>Onion</b>	Growth	Thrips  Purple blotch → 	<p><b>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</b></p> <p>Spray <b>Mencozeb</b> 25 gm and <b>hexaconazole</b> 15 ml in 10 liters of water at 10 days interval.</p>
<b>Brinjal &amp; Tomato</b>	Flowering/ fruiting	<b>Sucking pest</b> →	To control sucking pest spray <b>neem seed extract 5 %</b> or neem seed oil 50 ml and To control them

		& Fruit borer	spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
<b>Cabbage and cauliflower</b>	Growth & fruiting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.
<b>Chili</b>	Flowering & fruiting	Thrips, →  Wilting → Anthracnose → 	In present weather conditions severe attack of thrips has been reported. Spray 3 ml <b>spinosad</b> in 10 litres of water.  Drenching of Trichoderma viridi 4 gm in 1 liter of water.  Spray <b>Mencozeb 27</b> gm in 10 liters of water
<b>Mango</b>	Flowering	<b>Blossom midge and Leaf hopper</b> →   <b>Powdery Mildew</b> → 	To prevent damage spray <b>imidacloprid 5</b> ml/10 litres of water.      Spray 15 ml <b>Hexaconzole</b> in 10 liters of water.
<b>Livestock &amp; poultry</b> 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose. Put curtains on west side of poultry house.		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**





**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :94277 28523**



**Agro met Advisory Service bulletin No.-83 for the Amreli district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)


(Period 11-15 Jan.-2017)

**Weather forecast until 08-30 hrs of 18/01/2017**






**(3) AMRELI DISTRICT:**






	14/01/17	15/01/17	16/01/17	17/01/17	18/01/17
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	26	26	27	27	28
Min.Temp.(°C)	9	11	11	9	8
Total CC (octa)	0	1	3	4	1
RH-I (%)	32	45	48	48	45
RH-II (%)	22	27	28	31	25
Wind Speed (kmph)	16	9	17	20	20
Wind Direction(deg.)	56	139	261	76	76


**Agro meteorological Advisories:**

<b>General Advice</b>	<ul style="list-style-type: none"> <li>• <b>Rain fall:</b> Dry and clear to partly cloudy weather forecasted for next five days.</li> <li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 26-28°C and 8-11°C, respectively.</li> <li>• <b>Relative Humidity:</b> The RH is likely to be in the range of 18-45%</li> <li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be E, NE and 16-20 km/h, respectively.</li> <li>• <b>There is chance to further reduce in minimum temperature during 14-18 January, So to prevent cold/chilling injury irrigate field crops during evening time, carry out mulching and also make smoke in and around the field.</b></li> </ul>			
Crops	Stage	Insect/Pest/Variety	Advisory	
<b>Garlic</b>	Girthing	Thrips and  Alternaria leaf blight	<b>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</b>  The disease infestation observed in standing crop due to fog-moist weather. Increase periods between irrigations means do not over irrigate the crop. Make two Spray 27 gm Mencozeb/10 liters of water at 10 days interval.	



			
<b>Chick pea</b>	Flowering/ pod formation	Monitor the crop regularly for prodenia and heliethes infestation	“T” shaped bird perches in and around crop field to be installed for controlling insect population. If infestation cross ETL spray 3 ml <b>SPINOSAD</b> in 10 litres of water.
<b>Wheat</b>	Milking	-	Apply irrigation without stress.
<b>Cumin</b>	Flowering /grain formation	<b>Aphid and thrips</b>  <b>Blight and Powery mildew</b>    <b>Wilt</b>  	Under present weather condition attack of thrips and aphids has been reported spray <b>imidacloprid</b> <b>5 ml</b> /10 liters of water.  Under present weather condition spray <b>Hexaconazole</b> 15 ml/10 liters of water to prevent blight and powdery mildew infestation.    Stop irrigation to prevent further infection in standing crop.  
<b>Coriander</b>	Flowering & grain formation	Aphid, Thrips and Jassids	<b>Spray neem seed extract</b> 5% and <b>Thiomethoxam 5 gm/ 10 liters of water.</b> Under present weather condition spray <b>Hexaconazole</b> 15 ml/10 liters of water to prevent powdery mildew infestation.
<b>Onion</b>	Growth	Thrips   Purple blotch →	<b>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</b>  Spray <b>Mencozeb</b> 25 gm and <b>hexaconazole</b> 15 ml in 10 liters of water at 10 days interval.

			
<b>Brinjal &amp; Tomato</b>	Flowering/ fruiting	<b>Sucking pest</b> →  & <b>Fruit borer</b>	To control sucking pest spray <b>neem seed extract</b> 5 % or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
<b>Cabbage and cauliflower</b>	Growth & fruiting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.
<b>Chili</b>	Flowering & fruiting	Thrips, →  Wilting →  Anthracnose → 	In present weather conditions severe attack of thrips has been reported. Spray 3 ml <b>spinosad</b> in 10 litres of water.  Drenching of Trichoderma viridi 4 gm in 1 liter of water.  Spray <b>Mencozeb 27</b> gm in 10 liters of water
<b>Mango</b>	Flowering	<b>Blossom midge and Leaf hopper</b> →   <b>Powdery Mildew</b> → 	To prevent damage spray <b>imidacloprid</b> 5 ml/10 litres of water.         Spray 15 ml <b>Hexaconazole</b> in 10 liters of water.

<b>Livestock &amp; poultry</b> 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose. Put curtains on west side of poultry house.		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Gramin Krushi Mausam Seva**  
**Main Dry Farming Research Station**  
**Junagadh Agricultural University**  
**Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : (0281)2784722, Cell:94277 28523**



**Agro met Advisory Service bulletin No.-83 for the Surendranagar district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)



(Period 14-18Jan.-2017)






**Weather forecast until 08-30 hrs of 18/01/2017**






**(4) SURENDRANAGAR DISTRICT:**

	14/01/17	15/01/17	16/01/17	17/01/17	18/01/17
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	27	27	27	28	28
Min.Temp.(°C)	12	13	14	12	12
Total CC (octa)	0	2	3	5	0
RH-I (%)	27	41	44	50	48
RH-II (%)	15	18	22	26	23
Wind Speed (kmph)	16	11	18	16	17
Wind Direction(deg.)	80	76	319	42	50

**Agro meteorological advisories:**

<b>General Advice</b>	<ul style="list-style-type: none"> <li>• <b>Rain fall:</b> Dry and clear to partly cloudy weather forecasted for next five days.</li> <li>• <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 27-28°C and 12-14°C, respectively.</li> <li>• <b>Relative Humidity:</b> The RH is likely to be in the range of 15-50 %</li> <li>• <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be E, NW, NE and 11-18 km/h, respectively.</li> </ul>		
Crops	Stage	Insect/Pest/Variety	Advisory
<b>Garlic</b>	Githing	Thrips and  Alternaria leaf blight 	<p><b>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</b></p> <p>The disease infestation observed in standing crop due to fog-moist weather. Increase periods between irrigations means do not over irrigate the crop.            Make two Spray 27 gm Mencozeb/10 liters of water at 10 days interval.</p>
<b>Chick pea</b>	Flowering/ pod formation	Monitor the crop regularly for prodenia and	<p>“T” shaped bird perches in and around crop field to be installed for controlling insect population.            If infestation cross ETL spray 3 ml <b>SPINOSAD</b> in</p>

		heliothes infestation	10 litres of water.
<b>Wheat</b>	Grain formation	-	Apply irrigation without stress.
<b>Cumin</b>	Flowering /grain formation	<p><b>Aphid and thrips</b></p> <p><b>Blight and Powery mildew</b></p>  <p><b>Wilt</b></p> 	<p>Under present weather condition attack of thrips and aphids has been reported spray <b>imidacloprid 5 ml</b>/10 liters of water.</p> <p>Under present weather condition spray <b>Hexaconazole</b> 15 ml/10 liters of water to prevent blight and powdery mildew infestation.</p>  <p>Stop irrigation to prevent further infection in standing crop.</p> 
<b>Coriander</b>	Flowering & grain formation	Aphid, Thrips and Jassids	<p><b>Spray neem seed extract 5%</b> and <b>Thiomethoxam 5 gm/ 10 liters of water.</b></p> <p>Under present weather condition spray <b>Hexaconazole</b> 15 ml/10 liters of water to prevent powdery mildew infestation.</p>
<b>Onion</b>	Growth	<p>Thrips</p> <p>Purple blotch →</p> 	<p><b>For thrips control spray neem seed extract 5% and Make alternative sprays of profenofos 20 ml and lambda-cyhalothrin 14 ml in 10 liters of water at 10 days interval.</b></p> <p>Spray <b>Mencozeb</b> 25 gm and <b>hexaconazole</b> 15 ml in 10 liters of water at 10 days interval.</p>
<b>Brinjal &amp; Tomato</b>	Flowering/ fruiting	<b>Sucking pest</b> →	To control sucking pest spray <b>neem seed extract 5 %</b> or neem seed oil 50 ml and To control them spray acetamiprid 20 % SP@ 5.0 gm / 10 lit of

		& <b>Fruit borer</b>	water. In brinjal and tomato constant monitoring of borers by using pheromone traps 4-6 per acre is advised, If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against fruit borer.
<b>Cabbage and cauliflower</b>	Growth & fruiting	Diamond back moth	If insect population is above ETL than spraying of Spinosad 3.0 ml/10 litre of water is advised against diamond back moth in cauliflower and cabbage.
<b>Chili</b>	Flowering & fruiting	Thrips, →  Wilting → Anthracnose → 	In present weather conditions severe attack of thrips has been reported. Spray 3 ml <b>spinosad</b> in 10 litres of water.  Drenching of Trichoderma viridi 4 gm in 1 liter of water.  Spray <b>Mencozeb 27</b> gm in 10 liters of water
<b>Mango</b>	Flowering	<b>Blossom midge and Leaf hopper</b> →   <b>Powdery Mildew</b> → 	To prevent damage spray <b>imidacloprid 5</b> ml/10 litres of water.      Spray 15 ml <b>Hexaconzole</b> in 10 liters of water.
<b>Livestock &amp; poultry</b> 	Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Carry out vaccination for foot and mouth disease. Check up for reproductive diseases evaluation. Provide ad lib warm water to lactating animals. Keep young calves indoors during night time to avoid cold stress. Carry out sowing of maize for fodder purpose. Put curtains on west side of poultry house.		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**