



# હવामान आगाही आधारीत કૃષि સલાહ બુલેટીન

## ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ

તા. ૨૨-૦૪-૨૦૧૫ થી તા. ૨૬-૦૪-૨૦૧૫



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જી.કુ.યુ., તરઘડીયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિ હવામાન બુલેટીન નં: ૭/૨૦૧૫

પાછલા અઠવાડીયાનું રાજકોટનું હવામાન

હવામાન પરિબળો	ઉત્તર સૌરાષ્ટ્ર
વરસાદ (મી.મી)	0.00
મહત્તમ તાપમાન (સે.ગ્રે)	40-43.5
લઘુત્તમ તાપમાન (સે.ગ્રે)	23.2-25.3
ભેજ (%) સવાર	59-97
ભેજ (%) બપોરબાદ	24-33
પવનની ગતિ (કિમી/કલાક)	5.13-9.11

### હવામાન આગાહી: પત્રક-૫

ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા. ૨૨-૦૪-૨૦૧૫ થી તા. ૨૬-૦૪-૨૦૧૫

### કૃષિ સલાહ:

સામાન્ય

ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અતેના ગ્રામીણ કૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડીયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના.

**૧. રાજકોટ:** જીલ્લામાં આગામી તા. ૨૨-૦૪-૨૦૧૫ થી તા. ૨૬-૦૪-૨૦૧૫ દરમયાન હવામાન મુખ્યત્વે સુંકુ અને ચોખ્ખુ રહેશે. પવનની દિશા પણ્યમ અને વાયવ્ય રહેવાની અને પવનની ઝડપ ૧૫ થી ૧૮ ડિમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમયાન ૪૧ થી ૪૩ ડિગ્રી સેલ્સીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રિ દરમયાન ૨૫ થી ૨૭ ડિગ્રી સેલ્સીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ પેરામીટર	દિવસ-૧ ૨૨-૦૪-૧૫	દિવસ-૨ ૨૩-૦૪-૧૫	દિવસ-૩ ૨૪-૦૪-૧૫	દિવસ-૪ ૨૫-૦૪-૧૫	દિવસ-૫ ૨૬-૦૪-૧૫
વરસાદ (મી.મી)	0	0	0	0	0
મહત્તમ તાપમાન (સે.ગ્રે)	43	42	41	41	41
લઘુત્તમ તાપમાન (સે.ગ્રે)	26	26	26	25	25
વાદળની સ્થિતિ (ઓકટા)	0	0	0	0	0
મહત્તમ ભેજ (%) સવાર	41	54	66	76	81
લઘુત્તમ ભેજ (%) બપોરબાદ	16	17	20	27	35
પવનની ગતિ (કિમી/કલાક)	19	16	19	17	15
પવનની દિશા (ડિગ્રી)	300	270	270	270	280

- **૨. જામનગર** જીલ્લામાં આગામી તા.૨૨-૦૪-૨૦૧૫ થી તા.૨૬-૦૪-૨૦૧૫ દરમ્યાન હવામાન મુખ્યત્વે સુકુ અને ચોખ્યુ રહેશે. પવનની દિશા પણ્યમ રહેવાની અને પવનની ઝડપ ૧૫ થી ૨૦ ક્રિમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્વમાન દિવસ દરમ્યાન ૩૫ થી ૩૮ ડીગ્રી સેલ્ફીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રિ દરમ્યાન ૨૫ થી ૨૬ ડીગ્રી સેલ્ફીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ પેરામીટર	દિવસ-૧ ૨૨-૦૪-૧૫	દિવસ-૨ ૨૩-૦૪-૧૫	દિવસ-૩ ૨૪-૦૪-૧૫	દિવસ-૪ ૨૫-૦૪-૧૫	દિવસ-૫ ૨૬-૦૪-૧૫
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન તાપમાન (સે.ગ્રે)	36	36	37	38	38
લઘુત્તમ તાપમાન (સે.ગ્રે)	25	25	26	26	25
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0
મહત્વમાન ભેજ (%) સવાર	52	63	70	79	85
લઘુત્તમ ભેજ (%) બપોરબાદ	23	26	30	38	48
પવનની ગતિ (ક્રિમી/કલાક)	18	17	20	18	15
પવનની દિશા (ડીગ્રી)	280	260	270	260	260

- **અમરેલી** જીલ્લામાં આગામી તા.૨૨-૦૪-૨૦૧૫ થી તા.૨૬-૦૪-૨૦૧૫ દરમ્યાન હવામાન હવામાન મુખ્યત્વે સુકુ રહેવાની શક્યતા છે. પવનની દિશા પણ્યમ ની રહેવાની અને પવનની ઝડપ ૧૨ થી ૧૮ ક્રિમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્વમાન દિવસ દરમ્યાન ૪૦ થી ૪૧ ડીગ્રી સેલ્ફીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રિ દરમ્યાન ૨૫ થી ૨૬ ડીગ્રી સેલ્ફીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ પેરામીટર	દિવસ-૧ ૨૨-૦૪-૧૫	દિવસ-૨ ૨૩-૦૪-૧૫	દિવસ-૩ ૨૪-૦૪-૧૫	દિવસ-૪ ૨૫-૦૪-૧૫	દિવસ-૫ ૨૬-૦૪-૧૫
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન તાપમાન (સે.ગ્રે)	40	40	41	40	40
લઘુત્તમ તાપમાન (સે.ગ્રે)	25	26	26	25	25
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0
મહત્વમાન ભેજ (%) સવાર	42	48	60	69	72
લઘુત્તમ ભેજ (%) બપોરબાદ	21	21	22	27	34
પવનની ગતિ (ક્રિમી/કલાક)	18	13	17	15	12
પવનની દિશા (ડીગ્રી)	270	280	270	270	280

- **સુરેન્દ્રગઢ** જીલ્લામાં તા.૨૨-૦૪-૨૦૧૫ થી તા.૨૬-૦૪-૨૦૧૫ દરમ્યાન હવામાન મુખ્યત્વે સુકુ અને ચોખ્યુ રહેશે ત્થા તા. ૧૪ અને ૧૫ના રોજ પવનની દિશા પણ્યમની રહેવાની અને પવનની ઝડપ ૧૫ થી ૧૮ ક્રિમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્વમાન તાપમાન દિવસ દરમ્યાન ૪૦ થી ૪૨ ડીગ્રી સેલ્ફીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રિ દરમ્યાન ૨૫ થી ૨૭ ડીગ્રી સેલ્ફીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ પેરામીટર	દિવસ-૧ ૨૨-૦૪-૧૫	દિવસ-૨ ૨૩-૦૪-૧૫	દિવસ-૩ ૨૪-૦૪-૧૫	દિવસ-૪ ૨૫-૦૪-૧૫	દિવસ-૫ ૨૬-૦૪-૧૫
વરસાદ (મી.મી)	0	0	0	0	0
મહત્વમાન તાપમાન (સે.ગ્રે)	42	41	41	40	40
લઘુત્તમ તાપમાન (સે.ગ્રે)	26	27	26	26	26
વાદળની સ્થિતિ (ઓક્ટા)	0	0	0	0	0
મહત્વમાન ભેજ (%) સવાર	33	49	60	71	76
લઘુત્તમ ભેજ (%) બપોરબાદ	10	10	13	20	25
પવનની ગતિ (ક્રિમી/કલાક)	17	17	20	18	16
પવનની દિશા (ડીગ્રી)	280	260	260	260	270

- પાછલા અઠવાડિયા દરમિયાન વાદળાદી વરસાદી હવામાનના કારણે ઉનાળું મગફળીબ તલ અને મગના પાકમાં લીલી ઈયળ અને લશકરી ઈયળોનોઉપદ્રવ વધુ હોવાથી તેના નિયંત્રણ માટે ઈમામેકીટોન બેન્જોએટ ૭ ગ્રામ ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંઠવી)
ખાતરની ખાડ	પાક ખાતર પલટાવવું	—	<p>ઢોરના મળમુત્ર અને ખેત કચરામાંથી પૂર્ણ સરી ગયેલ ખાતરને ખાડમાંથી બણાર કાઢવું.</p>  
ઉનાળુ મગ	કુલ/શીંગ	સફેદ માખી, લીલી ઈયળ, તલના ભૂતીયા કુદાની ઈયળ	<ul style="list-style-type: none"> <li>➢ લીલોળીનું તેલ ૫૦ મીલી અથવા લીલોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી અને પ્રોફેનોફોસ ૨૦ મીલી ૧૦ લી પાણીમાં મિશ્ર કરી છંટકવ કરવો. જો ઈયળોનો વધુ ઉપદ્રવ હોય તો ઇન્ડોક્સાકાર્બ ૧૦ મીલી /૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકવ કરવો.</li>  </ul>
ઉનાળુ મગફળી	સુયા બેસવા/ડોડવા બંધાવા	લશકરી અને લીલી ઈયળ	<ul style="list-style-type: none"> <li>➢ કુલ/સુયા બેસવા અવસ્થાએ હળતુ પિયત આપવુ.</li> <li>➢ તેના નિયંત્રણ માટે એમામેકટીન બેન્જોએટ૭ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકવ કરવો.</li> </ul> 
ઉનાળુ તલ	કુલ/બેઢા બંધાવા	સફેદ માખી	<ul style="list-style-type: none"> <li>➢ તેના નિયંત્રણ માટે એસીટામાપ્રીડ ૫ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકવ કરવો.</li> </ul>

			<p>➢ તેમજ તલના પાકને કુલ અવસ્થાએ હળવુ પિયત આપવુ. ત્યારબાદ જમીનની પ્રત મુજબ ૧૦-૧૨ દિવસના ગણે પિયત આપવુ. વધારે પ્રમાણમાં ટૂંકા અંતરે પિયત આપવા નહિ, જરૂર કરતા વધારે ભેજથી તલમા સૂકારો અને કાળીયો રોગ આવવાની શક્યતા રહે છે.</p>
			
ચારાની જુવાર	વૃદ્ધિ	ગાભમારાની ઈયળ	<p>➢ તેના નિયંત્રણ માટે વેંડા સાયહેલોથીન ૫ ટકા ૧૦ મી.લી./૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકવ કરવો.</p>
રાગણી અને ટમેરી	કુલ/ફળ	શ્રીપસ, સફેદ માખી	<p>➢ લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી પાણીમા મીશ્ર કરી છંટકવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકવ કરવો.</p>
મરચી	કુલ/ફળ	શ્રીપસ	<p>પ્રોફેનોકોસ ૨૦ મી.લી. અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી પાણીમા મીશ્ર કરી છંટકવ કરવો</p>
ઝીડા	વૃદ્ધિ/કુલ/ શીંગ	શ્રીપસ, સફેદ માખી	<p>➢ તેના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી /૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકવ કરવો.</p>
વેલાવણા શાકભાજી	વૃદ્ધિ/ કુલ	<p>પાન કોરીયુ , શ્રીપસ, સફેદ માખી →</p> <p>તળધારો →</p>	<p>લીબોળીનો અર્ક ૫૦૦ મીલી અથવા લીબોળીનું તેલ ૫૦ મીલીનો છંટકવ ૧૫ દિવસના અંતરે ચાર વખત કરવો. વધુ ઉપદ્રવ હોય તો એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ /૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકવ કરવો.</p> <p>મેન્કોઝેક ૨૭ ગ્રામ અથવા કલોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકવ કરવા</p>
ગુવાર	વૃદ્ધિ/કુલ/શીંગ	શ્રીપસ, સફેદ માખી	<p>➢ તેના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી /૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકવ કરવો.</p>
તરબુચ	વૃદ્ધિ/કુલ	શ્રીપસ, સફેદ માખી	<p>➢ તેના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અને લીબોળીનું તેલ ૫૦ મીલી અથવા લીબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી /૧૦ લીટર પાણીમાં મીશ્ર કરી છંટકવ કરવો.</p>
આંબા	ફળ બંધાવા	<p>ફળમાખી</p> <p>પિયત</p>	<p>તેના નિયંત્રણ માટે મીથાઈલ યુજીનોંબ ટ્રેપનો ઉપયોગ કરી ફળમાખીનું નિયંત્રણ કરવુ</p> <p>આંબાના પાકમાં હળવુ પિયત આપવુ.</p>



પશુ/મરદા

પશુઓમાં ગળસુંઢો તથા ગાંઠીયા તાવના નિયંત્રણ માટે રસીકરણ કરવુ અને કરમીયાની ગોળી આપવી. દુધાળા પશુના આઉને ઝીક ઓકસાઈડ અથવા બોરીક પાવડર થી બરાબર સાફ કરવુ. તેમજ ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટાગ ૨ મી.લી. એક લીટર પાણીમાં નાખીને છાંટવી.

જા.નં.જૃકૃયુ/સંવે(સૂખે)/એગ્રોમેટ/બુલેટીન/ /૨૦૧૫  
તરઘડીયા( રાજકોટ), તારીખ: ૨૧/૦૪/૨૦૧૫ .

પ્રતિ,  
તંત્રીશ્રી/ ડાયરેક્ટરશ્રી,  
રાજકોટ.

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડિયોમાં ગામના ચોરાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિનામૂલ્યે પ્રસિદ્ધ કરવા નામ વિનંતી છે. સહકારની અપેક્ષા સહ આભાર.

સંશોધન વૈજ્ઞાનિક (સૂકી જેતી)  
જુનાગઢ કૃષિ યુનિવર્સિટી  
તરઘડીયા



**Main Dry Farming Research Station  
Junagadh Agricultural University  
Targhadia (Rajkot) - 360 003  
Phone: (0281)2784260(O), 2331686(R)  
Fax : ( 0281)2784722, Cell :94277 28523**



### **Agro met Advisory Service bulletin No.-7 for the Rajkot district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)  
(Period 22<sup>nd</sup> to 26<sup>th</sup> April- 2015)

#### **Significant past weather for the preceding week of Rajkot**

Rainfall (mm)	0.00
Maximum temperature (°C )	40-43.5
Minimum temperature (°C )	23.2-25.3
Relative Humidity (%) I	59-97
Relative Humidity (%) II	24-33
Wind speed (Kmph)	5.13-9.11

#### **Weather forecast until 08-30 hrs of 26/04/2015**

##### **(1) RAJKOT DISTRICT:**

	22/04/15	23/04/15	24/04/15	25/04/15	26/04/15
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	43	42	41	41	41
Min.Temp.(°C)	26	26	26	25	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	41	54	66	76	81
RH-II (%)	16	17	20	27	35
Wind Speed (kmph)	19	16	19	17	15
Wind Direction(deg.)	300	270	270	270	280

#### **Agro meteorological Advisories:**

<b>General Advice</b>	<ul style="list-style-type: none"> <li>➤ <b>Rain fall:</b> Dry and clear weather forecasted for the next.</li> <li>➤ <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 41-43 °C and 25-26 °C, respectively.</li> <li>➤ <b>Relative Humidity:</b> The RH is likely to be in the range of 16-81 %</li> <li>➤ <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W, NW and 15-19 km/h, respectively.</li> <li>➤ <b>Due to cloudy weather and off seasonal rainfall during last week heavy infestation of heliothis and prodenia observed in summer groundnut, sesame and green gram, to control these pest spray Emamectin Benzoate 7 gm in 10 liter of water.</b></li> </ul>		
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/Variety</b>	<b>Advisory</b>
<b>Summer green gram</b>	Flowering/pod formation	Heliothis and Catter pillar	Infestation of Heliothis and catter pillar observed

			To control the pest spray <b>profenofos</b> 50 % EC@ 20 ml /10 lit of water.
<b>Summer groundnut</b>	Pegging/pod formation	-	Apply light irrigation at flowering stage
<b>Summer Sesame</b>	Flowering/Capsule formation and development	White fly and prodenia	<b>Under present weather condition infestation of thrips, whiteflies and prodenia were observed in summer sown groundnut and sesame.</b> To control them spray <b>acetamiprid</b> 20 % SP@ 5.0 gm and emamectin benzoate 7 gm/10 lit of water.
<b>Fodder sorghum</b>	Growth	Shoot borer	Spray <i>lambda-cyhalothrin</i> 5 % EC, 10 ml/10 litre of water
<b>Lady finger</b>	Growing/fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Ridge gourd/ Bottle gourd/ Sponge gourd/ Bitter gourd</b>	Flowering/Fruiting	Leaf minor, Thrips, white fly  Downy mildew & Alternaria blight 	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) or <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.  Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
<b>Cluster bean</b>	Growing/fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Water melon</b>	Fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Brinjal and tomato</b>	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.  To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.

<b>Chili</b>	Flowering & fruiting	<b>Thrips</b> → 	In present weather conditions severe attack of thrips has been reported. Spray 20 ml <b>profenofos</b> in 10 litres of water.
<b>Mango</b>	Fruiting	<b>Blossom midge and Leaf hopper</b> → Fruit fly <b>Irrigation</b>	To prevent damage spray <b>imidacloprid 5</b> ml/10 litres of water.  Spray mixture of Fenthion 10 ml or DDVP 4 ml and Methyle ugenol 10 ml in 10 litres of water Apply light irrigation
<b>Livestock &amp; poultry</b>	Do Vaccination for <b>Haemorrhagic septicemia</b> (H.S.) disease and <b>Blue Quarter</b> (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Main Dry Farming Research Station  
Junagadh Agricultural University  
Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : ( 0281)2784722, Cell :94277 28523**



### **Agro met Advisory Service bulletin No.-7 for the Jamnagar district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 22<sup>nd</sup> to 26<sup>th</sup> April- 2015)

**Weather forecast until 08-30 hrs of 26/04/2015**

#### **(2) JAMNAGAR DISTRICT:**

	<b>22/04/15</b>	<b>23/04/15</b>	<b>24/04/15</b>	<b>25/04/15</b>	<b>26/04/15</b>
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	36	36	37	38	38
Min.Temp.(°C)	25	25	26	26	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	52	63	70	79	85
RH-II (%)	23	26	30	38	48
Wind Speed (kmph)	18	17	20	18	15
Wind Direction(deg.)	280	260	270	260	260

#### **Agro meteorological Advisories:-**

<b>General Advice</b>	<ul style="list-style-type: none"><li>➤ Rain fall Dry and mainly clear weather forecasted for the next five.</li><li>➤ <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 36-38 °C and 25-26 °C respectively.</li><li>➤ <b>Relative Humidity:</b> The RH is likely to be in the range of 23-85%</li><li>➤ <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W and 15-20 km/h, respectively.</li><li>➤ <b>Due to cloudy weather and off seasonal rainfall during last week heavy infestation of heliothis and prodenia observed in summer groundnut, sesame and green gram, to control these pest spray Emamectin Benzoate 7 gm in 10 liter of water.</b></li></ul>
-----------------------	---

Crops	Stage	Insect/Pest/Variety	Advisory
Summer green gram	Flowering	Heliothis and Catter pillar	Infestation of Heliothis and catter pillar observed To control the pest spray <b>profenofos</b> 50 % EC@ 20 ml /10 lit of water. 
Summer groundnut	Pegging/pod formation	-	Apply light irrigation at flowering stage 
Summer Sesame	Flowering/Capsule formation and development	White fly and prodenia	<b>Under present weather condition infestation of thrips, whiteflies and prodenia were observed in summer sown groundnut and sesame.</b> To control them spray <b>acetamiprid</b> 20 % SP@ 5.0 gm and emamectin benzoate 7 gm/10 lit of water. 
Fodder sorghum	Growth	Shoot borer	Spray <i>lambda-cyhalothrin</i> 5 % EC, 10 ml/10 litre of water
Lady finger	Growing/fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
Ridge gourd/Bottle gourd/	Flowering/Fruitting	Leaf minor, Thrips, white fly	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) or <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.

<b>Sponge gourd/ Bitter gourd</b>		Downy mildew & Alternaria blight 	Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
<b>Cluster bean</b>	Growing /fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Water melon</b>	Fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Brinjal and tomato</b>	Growing /fruiting	Sucking pest, Fruit and shoot borer	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.  To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
<b>Chili</b>	Flowering & fruiting	<b>Thrips</b> →  	In present weather conditions severe attack of thrips has been reported. Spray 20 ml <b>profenofos</b> in 10 litres of water.
<b>Mango</b>	Fruiting	<b>Blossom midge and Leaf hopper</b> → Fruit fly <b>Irrigation</b>	To prevent damage spray <b>imidacloprid 5</b> ml/10 litres of water.  Spray mixture of Fenthion 10 ml or DDVP 4 ml and Methyle ugenol 10 ml in 10 litres of water Apply light irrigation
<b>Livestock &amp; poultry</b> 	Do Vaccination for <b>Haemorrhagic septicemia</b> (H.S.) disease and <b>Blue Quarter</b> (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.		

**TechnicalOfficer, GKMS Unit, DFRS, JAU, Targhadia.**



**Main Dry Farming Research Station  
Junagadh Agricultural University  
Targhadia (Rajkot) - 360 003**

**Phone: (0281)2784260(O), 2331686(R)  
Fax : ( 0281)2784722, Cell :94277 28523**



**Agro met Advisory Service bulletin No.7 for the Amreli district**

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 22<sup>nd</sup> to 26<sup>th</sup> April- 2015)

**Weather forecast until 08-30 hrs of 26/04/2015**

**(3) AMRELI DISTRICT:**

	<b>22/04/15</b>	<b>23/04/15</b>	<b>24/04/15</b>	<b>25/04/15</b>	<b>26/04/15</b>
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	40	40	41	40	40
Min.Temp.(°C)	25	26	26	25	25
Total CC (octa)	0	0	0	0	0
RH-I (%)	42	48	60	69	72
RH-II (%)	21	21	22	27	34
Wind Speed (kmph)	18	13	17	15	12
Wind Direction(deg.)	270	280	270	270	280

### Agro meteorological Advisories:

<b>General Advice</b>	<ul style="list-style-type: none"> <li>➤ <b>Rain fall:</b> Dry and Clear weather forecasted for the next five.</li> <li>➤ <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 40-41°C and 25-26 °C, respectively.</li> <li>➤ <b>Relative Humidity:</b> The RH is likely to be in the range of 21-72%</li> <li>➤ <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W and 12-18 km/h, respectively.</li> <li>➤ <b>Due to cloudy weather and off seasonal rainfall during last week heavy infestation of heliothis and prodenia observed in summer groundnut, sesame and green gram, to control these pest spray Emamectin Benzoate 7 gm in 10 liter of water.</b></li> </ul>				
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/Variety</b>	<b>Advisory</b>		
<b>Summer green gram</b>	Flowering	Heliothis and Catter pillar	Infestation of Heliothis and catter pillar observed To control the pest spray <b>profenofos</b> 50 % EC@ 20 ml /10 lit of water. 		
<b>Summer groundnut</b>	Pegging/pod formation	-	Apply light irrigation at flowering stage 		
<b>Summer Sesame</b>	Flowering/Capsule formation and development	White fly and prodenia	<b>Under present weather condition infestation of thrips, whiteflies and prodenia were observed in summer sown groundnut and sesame.</b> To control them spray <b>acetamiprid</b> 20 % SP@ 5.0 gm and emamectin benzoate 7 gm/10 lit of water.		

			
<b>Fodder sorghum</b>	Growth	Shoot borer	Spray <i>lambda-cyhalothrin</i> 5 % EC, 10 ml/10 litre of water
<b>Lady finger</b>	Growing/fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Ridge gourd/Bottle gourd/Sponge gourd/Bitter gourd</b>	Flowering/Fruiting	Leaf minor, Thrips, white fly  Downy mildew & Alternaria blight 	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) or <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.  Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
<b>Cluster bean</b>	Growing/fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Water melon</b>	Fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Brinjal and tomato</b>	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.  To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
<b>Chili</b>	Flowering & fruiting	<b>Thrips</b> →	In present weather conditions severe attack of thrips has been reported. Spray 20 ml <b>profenofos</b> in 10 litres of water.

			
<b>Mango</b>	Fruiting	<b>Blossom midge and Leaf hopper</b> → Fruit fly <b>Irrigation</b>	To prevent damage spray <b>imidacloprid 5</b> ml/10 litres of water.  Spray mixture of Fenthion 10 ml or DDVP 4 ml and Methyle ugenol 10 ml in 10 litres of water Apply light irrigation
<b>Livestock &amp; poultry</b>	 Do Vaccination for <b>Haemorrhagic septicemia</b> (H.S.) disease and <b>Blue Quarter</b> (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**



**Main Dry Farming Research Station  
Junagadh Agricultural University  
Targhadia (Rajkot) - 360 003**  
**Phone: (0281)2784260(O), 2331686(R)**  
**Fax : (0281)2784722, Cell :94277 28523**



**Agro met Advisory Service bulletin No.-7 for the Surendranagar district**  
(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)  
(Period 22<sup>nd</sup> to 26<sup>th</sup> April- 2015)

**Weather forecast until 08-30 hrs of 26/04/2015**

**(4) SURENDRANAGAR DISTRICT:**

	22/04/15	23/04/15	24/04/15	25/04/15	26/04/15
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	42	41	41	40	40
Min.Temp.(°C)	26	27	26	26	26
Total CC (octa)	0	0	0	0	0
RH-I (%)	33	49	60	71	76
RH-II (%)	10	10	13	20	25
Wind Speed (kmph)	17	17	20	18	16
Wind Direction(deg.)	280	260	260	260	270

**Agro meteorological Advisories:**

<b>General Advice</b>	<ul style="list-style-type: none"><li>➤ <b>Rain fall:</b> Dry and clear weather forecasted for the next five days.</li><li>➤ <b>Temperature:</b> Maximum and minimum temperature likely to be in range of 40-42°C and 26-27 °C, respectively.</li><li>➤ <b>Relative Humidity:</b> The RH is likely to be in the range of 10-76 %</li><li>➤ <b>Wind Direction and Speed:</b> Direction and speed of wind is likely to be W and 16-18 km/h, respectively.</li><li>➤ <b>Due to cloudy weather and off seasonal rainfall during last week heavy infestation of heliothis and prodenia observed in summer groundnut, sesame and green gram, to control these pest spray Emamectin Benzoate 7 gm in 10 liter of water.</b></li></ul>		
<b>Crops</b>	<b>Stage</b>	<b>Insect/Pest/Var iety</b>	<b>Advisory</b>

<b>Summer green gram</b>	Flowering	Heliothis and Catter pillar	Infestation of Heliothis and catter pillar observed To control the pest spray <b>profenofos</b> 50 % EC@ 20 ml /10 lit of water. 
<b>Summer groundnut</b>	Pegging/pod formation	-	Apply light irrigation at flowering stage 
<b>Summer Sesame</b>	Flowering/Capsule formation and development	White fly and prodenia	<b>Under present weather condition infestation of thrips, whiteflies and prodenia were observed in summer sown groundnut and sesame.</b> To control them spray <b>acetamiprid</b> 20 % SP@ 5.0 gm and emamectin benzoate 7 gm/10 lit of water. 
<b>Fodder sorghum</b>	Growth	Shoot borer	Spray <i>lambda-cyhalothrin</i> 5 % EC, 10 ml/10 litre of water
<b>Lady finger</b>	Growing/fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Ridge gourd/ Bottle gourd/ Sponge gourd/</b>	Flowering/Fruiting	Leaf minor, Thrips, white fly	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) or <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval.

<b>Bitter gourd</b>		Downy mildew & Alternaria blight 	
<b>Cluster bean</b>	Growing/fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Water melon</b>	Fruiting	Thrips, white fly	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
<b>Brinjal and tomato</b>	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray <b>neem seed extract</b> 500 ml or neem seed oil 50 or neem oil 40 ml and <b>acetamiprid</b> 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
<b>Chili</b>	Flowering & fruiting	<b>Thrips</b> → 	In present weather conditions severe attack of thrips has been reported. Spray 20 ml <b>profenofos</b> in 10 litres of water.
<b>Mango</b>	Fruiting	<b>Blossom midge and Leaf hopper</b> → Fruit fly <b>Irrigation</b>	To prevent damage spray <b>imidacloprid 5</b> ml/10 litres of water.  Spray mixture of Fenthion 10 ml or DDVP 4 ml and Methyle ugenol 10 ml in 10 litres of water Apply light irrigation
<b>Livestock &amp; poultry</b> 	Do Vaccination for <b>Haemorrhagic septicemia</b> (H.S.) disease and <b>Blue Quarter</b> (B.Q.) in animals. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water.		

**Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.**