

ગ્રામીણ કૃષિ મૌસમ સેવા



હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન

ઉત્તર સૌરાષ્ટ્રખેત આબોહવાકીયવિભાગ

તા. ૨૫-૦૮-૨૦૧૭ થી તા. ૨૬-૦૮-૨૦૧૭



(ગ્રામીણ કૃષિ મૌસમ સેવા, સૂકી બેતી સંશોધન કેન્દ્ર, જુ.કૃ.યુ., તરધારીયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિહવામાન બુલેટીન નં. ૪૩/૨૦૧૭

પાછલા અઠવાડીયાનું રાજકોટનું હવામાન

હવામાન પરિબળો	
વરસાદ(મી.મી)	62.7
મહત્વમાન તાપમાન (સે.ગ્રે)	28.7-32.7
લઘુત્વમાન તાપમાન (સે.ગ્રે)	23.9-25.5
બેજ(%) સવાર	84-97
લેજ(%) બપોરખાદ	67-90
પવનની ગતિ (કિમી/કલાક)	5.0-7.7

હવામાન આગાહી પત્રક-૫

ઉત્તરસૌરાષ્ટ્રખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા. ૨૫-૦૮-૨૦૧૭ થી ૨૬-૦૮-૨૦૧૭

કૃષિ સલાહ:

સામાન્ય	<p>➢ ભારતસરકારશીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અતેના ગ્રામીણકૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરધારીયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના.</p> <p>રાજકોટ: જીવલામાં આગામી તા. ૨૫-૦૮-૨૦૧૭ થી તા. ૨૬-૦૮-૨૦૧૭ દરમ્યાન હવામાન લેજવાળું, ગરમ અને વાદળછાયું રહેશે તેમજ તા. ૨૬ થી ૨૮ ના રોજ હળવા અને તા. ૨૯ ના રોજ ભારે થી અતિ ભારે વરસાદ થવાની શક્યતા છે. પવનની દિશા પણ્યિમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૧૫ થી ૨૧ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્વમાન તાપમાન દિવસ દરમ્યાન ૩૧-૩૩ ડિગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્વમાન તાપમાન રાત્રી દરમ્યાન ૨૪-૨૫ ડિગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.</p>
---------	---

તારીખ	દિવસ-૧ ૨૫-૦૮-૧૭	દિવસ-૨ ૨૬-૦૮-૧૭	દિવસ-૩ ૨૭-૦૮-૧૭	દિવસ-૪ ૨૮-૦૮-૧૭	દિવસ-૫ ૨૯-૦૮-૧૭
પોરામીટર					
વરસાદ(મી.મી)	0	3	3	4	84
મહત્વમાન તાપમાન (સે.ગ્રે)	33	32	32	32	31
લઘુત્વમાન તાપમાન (સે.ગ્રે)	25	24	24	24	24

	વાદળનીસ્થિતિ(ઓકટા)	3	7	8	8
	મહતમલેજ(%) સવાર	86	82	82	84
	લધુતમલેજ (%) બપોરબાદ	64	65	66	65
	પવનનીગતિ(મી/કલાક)	17	21	18	15
	પવનનીદિશા(ડીગ્રી)	267	265	271	283

જમનગર: જીવ્લામાં આગામી તા. ૨૫-૦૮-૨૦૧૭ થી તા. ૨૯-૦૮-૨૦૧૭ દરમ્યાન હવામાન ભેજવાળું, ગરમ અને વાદળછાયું રહેશે. તેમજ તા. ૨૬ થી ૨૮ ના રોજ હળવા અને તા. ૨૯ ના રોજ ભારે વરસાદ થવાની શક્યતા છે. પવનની દિશા પણિયમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૧૮ થી ૨૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્વમાન તાપમાન દિવસ દરમ્યાન **૩૧-૩૩ ડીગ્રી સેલ્ફીયસ જેટલું** તેમજ લધુતમ તાપમાન રાત્રી દરમ્યાન **૨૫-૨૫ ડીગ્રી સેલ્ફીયસ જેટલું** રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પોરામીટર	૨૫-૦૮-૧૭	૨૬-૦૮-૧૭	૨૭-૦૮-૧૭	૨૮-૦૮-૧૭	૨૯-૦૮-૧૭
વરસાદ(મી.મી)	0	2	2	1	48
મહત્વમાન તાપમાન (સે.ગ્રે)	33	32	32	32	31
લધુત્વમાન તાપમાન (સે.ગ્રે)	25	25	25	25	25
વાદળનીસ્થિતિ(ઓકટા)	3	7	7	8	8
મહતમલેજ(%) સવાર	84	80	80	82	88
લધુતમલેજ(%) બપોરબાદ	67	68	68	67	74
પવનનીગતિ(કિમી/કલાક)	22	25	21	19	23
પવનનીદિશા(ડીગ્રી)	265	263	269	282	309

અમરેલી: જીવ્લામાં આગામી તા. ૨૫-૦૮-૨૦૧૭ થી તા. ૨૯-૦૮-૨૦૧૭ દરમ્યાન હવામાન ભેજવાળું, ગરમ અને વાદળછાયું રહેશે. તા. ૨૬ થી ૨૮ ના રોજ હળવા અને તા. ૨૯ ના રોજ ભારે થી અતિ ભારે વરસાદ થવાની શક્યતા છે. પવનની દિશા પણિયમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ **૧૪ થી ૧૮ કીમી/કલાક** રહેવાની શક્યતા છે. આ સમય ગાળામાં મહત્વમાન તાપમાન દિવસ દરમ્યાન **૩૧-૩૩ ડીગ્રી સેલ્ફીયસ જેટલું** તેમજ લધુતમ તાપમાન રાત્રી દરમ્યાન **૨૫-૨૫ ડીગ્રી સેલ્ફીયસ જેટલું** રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પોરામીટર	૨૫-૦૮-૧૭	૨૬-૦૮-૧૭	૨૭-૦૮-૧૭	૨૮-૦૮-૧૭	૨૯-૦૮-૧૭
વરસાદ(મી.મી)	0	5	5	4	84
મહત્વમાન તાપમાન (સે.ગ્રે)	33	33	32	32	31
લધુત્વમાન તાપમાન (સે.ગ્રે)	25	25	25	25	25
વાદળનીસ્થિતિ(ઓકટા)	5	7	8	8	8

	મહતમલેજ(%) સવાર	86	82	83	84	91
	લધુતમલેજ(%) બપોરબાદ	64	66	66	65	73
	પવનનીગતિક્રિ/કલાક)	17	19	16	14	16
	પવનનીદિશા(ડીગ્રી)	270	267	274	288	316

સુરેન્દ્રનગર જલ્લામાં આગામી તા. ૨૫-૦૮-૨૦૧૭ થી તા. ૨૯-૦૮-૨૦૧૭ દરમાન હવામાન બેજવાળું, ગરમ અને વાદળાયું રહેશે તા. ૨૬ થી ૨૮ ના રોજ હળવા અને તા. ૨૮ ના રોજ ભારે વરસાદ થવાની શક્યતા છે. પવનની દિશા નૈऋત્ય અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૧૪ થી ૧૮ કીમી/કલાક રહેવાની શક્યતા છે. આ સમય ગાળામાં મહતમ તાપમાન દિવસ દરમાન ૩૧-૩૩ ડીગ્રી સેલ્ફીયસ જેટલું તેમજ લધુતમ તાપમાન રાત્રી દરમાન ૨૪-૨૫ ડીગ્રી સેલ્ફીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૨૫-૦૮-૧૭	૨૬-૦૮-૧૭	૨૭-૦૮-૧૭	૨૮-૦૮-૧૭	૨૯-૦૮-૧૭
વરસાદ(મી.મી)	0	2	2	2	50
મહતમતાપમાન (સ.ગ્રે)	33	32	32	32	31
લધુતમતાપમાન (સ.ગ્રે)	25	24	24	24	24
વાદળનીસ્થિતિ(ઓકટા)	4	7	7	8	8
મહતમલેજ(%) સવાર	80	77	77	80	87
લધુતમલેજ(%) બપોરબાદ	55	56	57	59	64
પવનનીગતિક્રિ/કલાક)	19	19	16	14	14
પવનનીદિશા(ડીગ્રી)	263	266	270	284	312

૨૮ ઓગસ્ટ સુધી દરમાન હવામાન બેજવાળું, ગરમ અને વાદળાયું રહેશે. તેમજ રાજકોટ અને અમરેલી જલ્લામાં તા. ૨૬ થી ૨૮ ના રોજ હળવા અને તા. ૨૮ ના રોજ ભારે થી અતિ ભારે વરસાદ થવાની શક્યતા છે જામનગર અને સુરેન્દ્રનગર જલ્લામાં તા. ૨૬ થી ૨૮ ના રોજ હળવા અને તા. ૨૮ ના રોજ ભારે થી અતિ ભારે વરસાદ થવાની શક્યતા છે. પવનની દિશા વાયવ્ય અને પણ્યમની રહેવાની અને પવનની ઝડપ ૧૪ થી ૨૫ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહતમ તાપમાન દિવસ દરમાન ૩૧-૩૩ ડીગ્રી સેલ્ફીયસ જેટલું તેમજ લધુતમ તાપમાન રાત્રી દરમાન ૨૪-૨૫ ડીગ્રી સેલ્ફીયસ જેટલું રહેવાની સંભાવના છે.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાતાજત	કૃપિ સલાહ (૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
મગફળી	કૂલ અને સુયા	સફેદ મુંડા 	ફોરેટ ૧૦ ટકા દાઢાદાર દવા ૨૫ કી.ગ્રા. પ્રતિ હેક્ટરે જમીનમાં પુરતો ભેજ હોય ત્યારે નાખવી અથવા કલોરોપાયરીઝેસ દવા ૪ લીટર પ્રતિ હેક્ટરે પિયત પાણી સાથે અથવા કલોરોપાયરીઝેસ દવા ૨૫ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી જમીનમાં હુલની ફરતે રેડવું. પાનના ટપકાના રોગના આગોતરા નિયંત્રણ માટે

		ચીતરી અને ગેડુ 	હેક્ઝાકોનાજોલ ૧૫ મિલી અથવા ટેલ્યુકોનાજોલ ૧૦ મિલી ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો.
કપાસ	ચાપવા	ગુલાબી ઈયળ, લીલા તડતડીયા/શ્રીપ્સ	કપાસમાં ગુલાબી ઈયળના નર કુદાને એકત્રીત કરવા માટે હેક્ટરે ૪૦ ની સંખ્યામાં ફેરોમેન ટ્રેપ ગોઠવવાથી ઈયળોનું શરૂઆતની અવસ્થાથી જ નિયત્રણ કરી શકાય . કપાસમાં ગુલાબી ઈયળથી નુકશાન પામેલ ફૂલ અને ચાપવા તોડીને જમીનમાં દાટી દેવા. વધુ નુકશાન હોય તો પ્રોફેનોફોસ ૫૦% ઇ.સી. ૨૦ મી.લી. અથવા સાયપરમેથીન ૧૦ મિલી બીવેરીયા બાસીઆના ફુગનો પાઉડર ૪૦ ગ્રામ ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો. લીલા તડતડીયા/શ્રીપ્સના નિયત્રણ માટે ઇમીડાંકલોપ્રીડ ૧૭.૮ ટકા ઇસી ૭ મિલી ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો.
તલ	ફૂલ	પાનવાળનાર(માથા બાંધનાર) ઈયળ 	કવીનાલફોસ ૨૦ મિલી અથવા કાટેપ હાયફ્રોક્લોરાઇડ ૧૫ ગ્રામ ૧૦ લીટર પાણીમા મિશ્ર કરી છંટકાવ કરવો.
બાજરી	ફૂલ	-	-
કઠોળ પાકો	શીગમાં દાણાભરાવા	-	પાકેલી શીંગો ઉતારી લેવી
શાકભાજુ	રોપણી/ વાવેતર	-	ઘરું તૈયાર હોય તો મરચી, ટમેટી, રીંગણ અને કોબીજુની ફેર રોપણી કરવી
સીડા	વૃદ્ધિ/ફૂલ/શીંગ	શ્રીપ્સ, સફેદ માખી	➢ તેના નીયત્રણ માટે એસીટામાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અને લીંબોળીનું તેલ ૫૦ મિલી અથવા લીંબોળીના મીજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
તુરીયા, ફુધી, ગાલકા, કાકડી અને	વૃદ્ધિ/ફૂલ	પાન કોરીયું , શ્રીપ્સ, સફેદ માખી	➢ લીંબોળીનો અર્ક ૫૦૦ મિલી અથવા લીંબોળીનું તેલ ૫૦ મિલીનો છંટકાવ ૧૫ દિવસના અંતરે ચાર વખત કરવો. વધુ ઉપક્રમ હોય તો એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ / ૧૦ લીટર

કારેલા		તલધારો	<p>પાણીમા મિશ્ર કરી છંટકાવ કરવો.</p> <ul style="list-style-type: none"> ➤ મેન્ડોઝેબ ૨૭ ગ્રામ અથવા કલોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઇલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.
ગુવાર	વૃધ્ઘા/કુલ/શીંગા	શ્રીપ્સ, સફેદ માખી	<ul style="list-style-type: none"> ➤ તેના નિયંત્રણ માટે એસીટામાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
રીંગણી અને ટમેટી	કુલ/ફળ	શ્રીપ્સ, સફેદ માખી, ફળ અને ડોકા કોરી ખાનાર ઈચ્છા	<ul style="list-style-type: none"> ➤ લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળી ના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપક્રમ હોય તો એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. રીંગણી અને ટમેટીમાં ફળ અને ડોકા કોરી ખાનાર ઈચ્છાનો ઉપક્રમ વધુ હોય તો સ્પીનોસાડ ૩ મીલી /૧૦ લીટર પાણીમાં ઓગાળી છંટકાવ કરવો.
મરચી	કુલ/ફળ	શ્રીપ્સ	<ul style="list-style-type: none"> ➤ પ્રોફેનોફ્રોસ ૨૦ મીલી અને લીંબોળી નું તેલ ૫૦ મીલી અથવા લીંબોળી ના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
લીંગુ	-	-	સતત બેજવાળા હવામાનના કારણે પાનાં ટપકા અને ગુંદરીયા રોગનો ઉપક્રમ નાં થાય તે માટે કોપર ઓક્ષિક્લોરાઇડ ૪૦ ગ્રામ અને સ્ટ્રેપટોસાયક્લીન ૧.૫ ગ્રામ ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
પશુ/મરધા		દુધાળા પશુના આઉને ઝીંક ઓક્સાઇડ અથવા બોરીક પાવડર થી બરાબર સાફ કરવું. તેમજ ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથીન અથવા એમીટ્રાલ ૨ મી.લી. એક લીટર પાણીમાં નાખીને છાંટવી અને કરમીયાની ગોળી આપવી તેમજ ટીબી અને જે.ડી.ના રોગની પરખ કરાવવી તેમજ બુસેલ્તાનું રસીકરણ કરાવવું. વરસાદના સમયમાં પશુઓને સુરક્ષિત રાખવા. ચોમાસા દરમ્યાન પશુઓના રહેઠાણની જગ્યાની વ્યવસ્થિત સફાઈ રાખવી અને માખીઓના ઉપક્રમથી બચવા ફિનાઇલ છાંટવું. ખુલ્લામાં રાખેલ ધાંસચારો વરસાદથી પલળો/બગાડ નાહિ તે માટે જરૂરી પગલા ભરવા.	

જુહૂ/ટેકનિકલ ઓફિસર.(સુઅ.)/એગ્રોમેટ/બુલેટીન/

/૨૦૧૭,

તરધારા(રાજકોટ), તારીખ: ૨૪/૦૮/૨૦૧૭.

પ્રતિ,

તંત્રીશ્રી/ડાયેક્ટરશ્રી,

રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં , રેડિયોમાં ગામના ચો રાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમ માં
વિનામૂલ્યેપ્રસિધ્ધ કરાવવા નમ્ર વિનતી છે. સહકારનીઅપેક્ષા સહ આભાર.

ટેકનીકલ ઓફિસર
સુકી ખેતીસંશોધન કેન્દ્ર
જુનાગઢ કૃષિ યુનિવર્સિટી
તરધારા



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-43 for the Rajkot district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 25 to 29-Aug.- 2017)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	62.7
Maximum temperature (°C)	28.7-32.7
Minimum temperature (°C)	23.9-25.5
Relative Humidity (%) I	84-97
Relative Humidity (%) II	67-90
Wind speed (Kmph)	5.0-7.7

Weather forecast until 08-30 hrs of 29/08/2017

(1) RAJKOT DISTRICT:

	25/08/17	26/08/17	27/08/17	28/08/17	29/08/17
Rainfall (mm)	0	3	3	4	84
Max.Temp.(°C)	33	32	32	32	31
Min.Temp.(°C)	25	24	24	24	24
Total CC (octa)	3	7	8	8	8
RH-I (%)	86	82	82	84	91
RH-II (%)	64	65	66	65	73
Wind Speed (kmph)	17	21	18	15	18
Wind Direction(deg.)	267	265	271	283	319

Agro meteorological Advisories:

General Advice	<ul style="list-style-type: none">Rain fall: Humid, Warm and generally cloudy to cloudy weather forecasted for next five days. light rainfall during 26 to 28 & Heavy to very heavy rainfall during 29 August forecasted.Temperature: Maximum and minimum temperature likely to be in range of 31-33°C and 24-25°C, respectively.Relative Humidity: The RH at morning and noon is likely to be in the range of 82-91 % and 64-73 %, respectively.Wind Direction and Speed: Direction and speed of wind is likely to be W, NW and
-----------------------	---

	<p>15-21 km/h, respectively.</p> <ul style="list-style-type: none"> Carry out interculturing, weeding in all crops for aeration, removal of weeds and conservation of soil moisture. 		
Crops	Stage	Insect/Pest/Variety	Advisory
Kharif groundnut	Flowering, pegging	White grub  Tikka & rust 	<p>Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha.</p> <p>Spray Tebuconazole 10 ml/10 lit of water to prevent leaf spot infestation & rust diseases infestation.</p>
Cotton	Squaring	Thrips, jassids & aphid Pink bollworm	Under present weather infestation of thrips, jassids & aphid observed in cotton, spray Imidacloprid 17.8 % 5 ml or Thiomethoxam 3 gram or Dimethoate 10 ml in 10 liters of water, if population is above ETL. For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha. <p>Farmers are advised to pick and bury the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 ml in 10 liters of water.</p>
Sesame	Flowering	Leaf webber 	Spray Cartep Hydrochloride 25 gram ml in 10 litres of water.
Castor	Growing/flowering	Semi looper & prodenia→ 	Heavy infestation of semilooper observed in castor, so spray Chlorpyro phos 30 ml in 10 liters of water.
Pearl millet	Flowering initiation	-	-
Pulses	Pod formation & development	-	Pick mature pod
Fodder Sorghum	Growing	Sorghum-Gundari, GFS-4 & 5, Maize	-

& Maize		African Tall	
Vegetables	Transplanting	Farmers are advised	Transplanting of brinjal, chilli and early cauliflower
Lady finger	Flowering/fruiting	White fly,	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
Ridge gourd/ Bottle gourd/ Sponge gourd/ Bitter gourd/ Cucumbers	Flowering/Fruiting	White fly Downy mildew & Alternaria blight 	Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Due to continuous humid weather Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Brinjal and tomato	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips →  	In present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofos in 10 litres of water.
Lemon	-	Citrus canker	Due to continuous humid weather spray Copper oxychloride 40 gram & 1.5 gram Streptocycline for prevention of citrus canker & gummosis disease
Livestock & poultry 		Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Do Brucellosis calfwood vaccination. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Do Arrangement for artificial light to birds to maintain egg production in poultry farm.	

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-43 for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 25 to 29-Aug.- 2017)

Weather forecast until 08-30 hrs of 29/08/2017

(2) JAMNAGAR DISTRICT:

	25/08/17	26/08/17	27/08/17	28/08/17	29/08/17
Rainfall (mm)	0	2	2	1	48
Max.Temp.(°C)	33	32	32	32	31
Min.Temp.(°C)	25	25	25	25	25
Total CC (octa)	3	7	7	8	8
RH-I (%)	84	80	80	82	88
RH-II (%)	67	68	68	67	74
Wind Speed (kmph)	22	25	21	19	23
Wind Direction(deg.)	265	263	269	282	309

Agro meteorological Advisories:-

General Advice	<ul style="list-style-type: none">Rain fall: Humid, Warm and generally cloudy to cloudy weather forecasted for next five days. light rainfall during 26 to 28 & Heavy rainfall during 29 August forecastedTemperature: Maximum and minimum temperature likely to be in range of 31-33°C and 25-25°C, respectively.Relative Humidity: The RH at morning and noon is likely to be in the range of 80-88 % and 67-74 %, respectively.Wind Direction and Speed: Direction and speed of wind is likely to be W, NW and 19-25 km/h, respectively.Carry out interculturing, weeding in all crops for aeration, removal of weeds and conservation of soil moisture.
-----------------------	--

Crops	Stage	Insect/Pest/Variety	Advisory
Kharif groundnut	Flowering, pegging	White grub 	Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha.

		Tikka & rust 	Spray Tebuconazole 10 ml/10 lit of water to prevent leaf spot infestation & rust diseases infestation.
Cotton	Squaring	Thrips, jassids & aphid Pink bollworm	Under present weather infestation of thrips, jassids & aphid observed in cotton, spray Imidacloprid 17.8 % 5 ml or Thiomethoxam 3 gram or Dimethoate 10 ml in 10 liters of water, if population is above ETL. For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha. Farmers are advised to pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 mi in 10 liters of water.
Sesame	Flowering	Leaf webber 	Spray Cartep Hydrochloride 25 gram ml in 10 litres of water.
Castor	Growing/flowering	Semi looper & prodenia→ 	Heavy infestation of semilooper observed in castor, so spray Chlorpyro phos 30 ml in 10 liters of water.
Pearl millet	Flowering initiation	-	-
Pulses	Pod formation & developmet	-	Pick mature pod
Fodder Sorghum & Maize	Growing	Sorghum-Gundari, GFS-4 & 5, Maize African Tall	-
Vegetables	Transplanting	Farmers are advised Transplanting of brinjal, chilli and early cauliflower	
Lady finger	Flowering/fruiting	White fly,	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
Ridge gourd/ Bottle gourd/	Flowering/Fruiting	White fly	Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.

Sponge gourd/ Bitter gourd/ Cucumbers		Downy mildew & Alternaria blight 	Due to continuous humid weather Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Brinjal and tomato	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips →  	In present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofos in 10 litres of water.
Lemon	-	Citrus canker	Due to continuous humid weather spray Copper oxychloride 40 gram & 1.5 gram Streptocycline for prevention of citrus canker & gummosis disease
Livestock & poultry 		Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Do Brucellosis calfwood vaccination. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Do Arrangement for artificial light to birds to maintain egg production in poultry farm.	

TechnicalOfficer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-43 for the Amreli district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 25 to 29-Aug.- 2017)

Weather forecast until 08-30 hrs of 29/08/2017

(3) AMRELI DISTRICT:

	25/08/17	26/08/17	27/08/17	28/08/17	29/08/17
Rainfall (mm)	0	5	5	4	84
Max.Temp.(°C)	33	33	32	32	31
Min.Temp.(°C)	25	25	25	25	25
Total CC (octa)	5	7	8	8	8
RH-I (%)	86	82	83	84	91
RH-II (%)	64	66	66	65	73
Wind Speed (kmph)	17	19	16	14	16
Wind Direction(deg.)	270	267	274	288	316

Agro meteorological Advisories :

General Advice	<ul style="list-style-type: none"> • Rain fall: Humid, Warm and generally cloudy to cloudy weather forecasted for next five days. light rainfall during 26 to 28 & Heavy to very heavy rainfall during 29 August forecasted • Temperature: Maximum and minimum temperature likely to be in range of 31-33°C and 25-25°C, respectively. • Relative Humidity: The RH at morning and noon is likely to be in the range of 82-91 % and 64-73 %, respectively. • Wind Direction and Speed: Direction and speed of wind is likely to be W, NW and 14-19 km/h, respectively. • Carry out interculturing, weeding in all crops for aeration, removal of weeds and conservation of soil moisture.
-----------------------	--

Crops	Stage	Insect/Pest/Variety	Advisory
Kharif groundnut	Flowering, pegging	White grub 	Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha.

		Tikka & rust 	Spray Tebuconazole 10 ml/10 lit of water to prevent leaf spot infestation & rust diseases infestation.
Cotton	Squaring	Thrips, jassids & aphid Pink bollworm	Under present weather infestation of thrips, jassids & aphid observed in cotton, spray Imidacloprid 17.8 % 5 ml or Thiomethoxam 3 gram or Dimethoate 10 ml in 10 liters of water, if population is above ETL. For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha. Farmers are advised to pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 mi in 10 liters of water.
Sesame	Flowering	Leaf webber 	Spray Cartep Hydrochloride 25 gram ml in 10 litres of water.
Castor	Growing/	Semi looper & prodenia→ 	Heavy infestation of semilooper observed in castor, so spray Chlorpyro phos 30 ml in 10 liters of water.
Pearl millet	Flowering initiation	-	-
Pulses	Pod formation & development	-	Pick mature pod
Fodder Sorghum & Maize	Growing	Sorghum-Gundari, GFS-4 & 5, Maize African Tall	-
Vegetables	Transplanting	Farmers are advised Transplanting of brinjal, chilli and early cauliflower	
Lady finger	Flowering/fruiting	White fly,	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
Ridge gourd/ Bottle	Flowering/Fruiting	White fly	Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP@ 5.0 gm / 10 lit

gourd/ Sponge gourd/ Bitter gourd/ Cucumbers		Downy mildew & Alternaria blight 	of water at 15 days interval. Due to continuous humid weather Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Brinjal and tomato	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips →  	In present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofosin 10 litres of water.
Lemon	-	Citrus canker	Due to continuous humid weather spray Copper oxychloride 40 gram & 1.5 gram Streptocycline for prevention of citrus canker & gummosis disease
Livestock & poultry 		Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Do Brucellosis calfwood vaccination. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Do Arrangement for artificial light to birds to maintain egg production in poultry farm.	

TechnicalOfficer, GKMS Unit, DFRS, JAU, Targhadia.



GraminKrushiMausamSeva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell:94277 28523



Agro met Advisory Service bulletin No.-43 for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 25 to 29-Aug.- 2017)

Weather forecast until 08-30 hrs of 29/08/2017

(4) SURENDRANAGAR DISTRICT:

	25/08/17	26/08/17	27/08/17	28/08/17	29/08/17
Rainfall (mm)	0	2	2	2	50
Max.Temp.(°C)	33	32	32	32	31
Min.Temp.(°C)	25	24	24	24	24
Total CC (octa)	4	7	7	8	8
RH-I (%)	80	77	77	80	87
RH-II (%)	55	56	57	59	64
Wind Speed (kmph)	19	19	16	14	14
Wind Direction(deg.)	263	266	270	284	312

Agro meteorological advisories:

General Advice	<ul style="list-style-type: none">Rain fall: Humid, Warm and generally cloudy to cloudy weather forecasted for next five days. light rainfall during 26 to 28 & Heavy rainfall during 29 August forecastedTemperature: Maximum and minimum temperature likely to be in range of 31-33°C and 24-25°C, respectively.Relative Humidity: The RH at morning and noon is likely to be in the range of 77-87 % and 55-64 %, respectively.Wind Direction and Speed: Direction and speed of wind is likely to be W, NW and 14-19 km/h, respectively.Carry out interculturing, weeding in all crops for aeration, removal of weeds and conservation of soil moisture.
-----------------------	--

Crops	Stage	Insect/Pest/Variety	Advisory
Kharif groundnut	Flowering, pegging	White grub 	Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha.

		Tikka & rust 	Spray Tebuconazole 10 ml/10 lit of water to prevent leaf spot infestation & rust diseases infestation.
Cotton	Squaring	Thrips, jassids & aphid Pink bollworm	Under present weather infestation of thrips, jassids & aphid observed in cotton, spray Imidacloprid 17.8 % 5 ml or Thiomethoxam 3 gram or Dimethoate 10 ml in 10 liters of water, if population is above ETL. For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 6-8/ha. Farmers are advised to pick and burry the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml or cypermethrin 10 mi in 10 liters of water.
Sesame	Flowering	Leaf webber 	Spray Cartep Hydrochloride 25 gram ml in 10 litres of water.
Castor	Growing	Semi looper & prodenia→ 	Heavy infestation of semilooper observed in castor, so spray Chlorpyro phos 30 ml in 10 liters of water.
Pearl millet	Flowering initiation	-	-
Pulses	Pod formation & development	-	Pick mature pod
Fodder Sorghum & Maize	Growing	Sorghum-Gundari, GFS-4 & 5, Maize African Tall	-
Vegetables	Transplanting	Farmers are advised Transplanting of brinjal, chilli and early cauliflower	
Lady finger	Flowering/fruiting	White fly,	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval.
Ridge gourd/ Bottle	Flowering/Fruiting	White fly	Sparyneem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP@ 5.0 gm / 10 lit

gourd/ Sponge gourd/ Bitter gourd/ Cucumbers		Downy mildew & Alternaria blight 	of water at 15 days interval. Due to continuous humid weather Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Brinjal and tomato	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips →  	In present weather conditions severe attack of thrips has been reported. Spray 20 ml profenofosin 10 litres of water.
Lemon	-	Citrus canker	Due to continuous humid weather spray Copper oxychloride 40 gram & 1.5 gram Streptocycline for prevention of citrus canker & gummosis disease
Livestock & poultry 		Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Do Brucellosis calfwood vaccination. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Due to chance of rainfall keep animals indoors during rainfall. Keep fodder in safe place to avoid spoilage due to rainfall. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Do Arrangement for artificial light to birds to maintain egg production in poultry farm.	

TechnicalOfficer, GKMS Unit, DFRS, JAU, Targhadia.