



ગ્રામીણ કૃષિ મૌસમ સેવા

હવામાન આગાહી આધારીત કૃષિસલાહ બુલેટીન

ઉત્તર સૌરાષ્ટ્રખેત આબોહવાકીય વિભાગ

તા.૨૮-૦૯-૨૦૧૬ થી તા.૦૨-૧૦-૨૦૧૬



(ગ્રામીણ કૃષિ મૌસમસેવા, સૂકી ખેતી સંશોધન કેન્દ્ર, જ.કૃ.યુ., તરઘડિયા અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

કૃષિહવામાન બુલેટીન નં. ૫૨/૨૦૧૬

પાછલા અઠવાડિયાનું રાજકોટનું હવામાન

હવામાન પરિબલો	
વરસાદ(મી.મી.)	0.0
મહત્તમ તાપમાન (સે.ગ્રે.)	32.0-32.8
લઘુત્તમ તાપમાન (સે.ગ્રે.)	21.5-23.8
ભેજ(%) સવાર	84-89
ભેજ(%) બપોરબાદ	58-64
પવનની ગતિ (કિમી/કલાક)	6.8-8.1

હવામાન આગાહી: પત્રક-ક

ઉત્તર સૌરાષ્ટ્રખેત આબોહવાકીય વિભાગ-હવામાન આગાહી

તા.૨૮-૦૯-૨૦૧૬ થી તા.૦૨-૧૦-૨૦૧૬

કૃષિ સલાહ:

સામાન્ય	➤ ભારત સરકારશ્રીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણ કૃષિ મૌસમ સેવા વિભાગ, જુનાગઢ કૃષિ યુનિવર્સિટી, તરઘડિયા તરફથી જણાવવામાં આવે છે કે, ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના રાજકોટ: જીલ્લામાં આગામી તા.૨૮-૦૯-૨૦૧૬ થી તા.૦૨-૧૦-૨૦૧૬ દરમિયાન હવામાન ભેજવાળું અને અંશતઃ વાદળછાયું રહેશે અને તા.૨ ઓક્ટોબરના રોજ હળવા વરસાદની શક્યતા છે. પવનની દિશા પશ્ચિમની રહેવાની અને પવનની ઝડપ ૧૨ થી ૨૦ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૩-૩૩ ડીગ્રી સેલ્શિયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૨-૨૩ ડીગ્રી સેલ્શિયસ જેટલું રહેવાની સંભાવના છે.					
	તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
	પેરામીટર	૨૮-૦૯-૧૬	૨૯-૦૯-૧૬	૩૦-૦૯-૧૬	૦૧-૧૦-૧૬	૦૨-૧૦-૧૬
	વરસાદ(મી.મી.)	0	0	0	0	10
	મહત્તમ તાપમાન (સે.ગ્રે.)	33	33	33	33	33
	લઘુત્તમ તાપમાન (સે.ગ્રે.)	23	22	22	22	22
	વાદળની સ્થિતિ(ઓકટા)	2	1	1	2	2

મહત્તમ ભેજ (%) સવાર	90	91	90	89	83
લઘુત્તમ ભેજ (%) બપોરબાદ	53	50	51	49	49
પવનની ગતિ (કિમી/કલાક)	18	18	15	12	20
પવનની દિશા (ડીગ્રી)	282	276	276	286	287

જામનગર: જીલ્લામાં આગામી તા. ૨૪-૦૯-૨૦૧૬ થી તા. ૨૮-૦૯-૨૦૧૬ દરમિયાન હવામાન ભેજવાળું અને અંશતઃ વાદળાળું રહેશે. વરસાદની શક્યતા નથી. પવનની દિશા પશ્ચિમની રહેવાની અને પવનની ઝડપ ૧૧ થી ૨૦ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૨-૩૨ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૪-૨૪ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૨૮-૦૯-૧૬	૨૯-૦૯-૧૬	૩૦-૦૯-૧૬	૦૧-૧૦-૧૬	૦૨-૧૦-૧૬
વરસાદ(મી.મી)	0	0	0	0	0
મહત્તમ તાપમાન (સે.ગ્રે)	32	32	32	32	32
લઘુત્તમ તાપમાન (સે.ગ્રે)	24	24	24	24	24
વાદળની સ્થિતિ(ઓકટા)	2	1	0	2	1
મહત્તમ ભેજ (%) સવાર	89	89	88	88	83
લઘુત્તમ ભેજ (%) બપોરબાદ	57	55	56	55	55
પવનની ગતિ (કિમી/કલાક)	18	19	15	11	20
પવનની દિશા (ડીગ્રી)	276	270	268	280	285

અમરેલી: જીલ્લામાં આગામી તા. ૨૮-૦૯-૨૦૧૬ થી તા. ૦૨-૧૦-૨૦૧૬ દરમિયાન હવામાન ભેજવાળું અને અંશતઃ વાદળાળું રહેશે અને તા.૨ ઓક્ટોબરના રોજ હળવા વરસાદની શક્યતા છે. પવનની દિશા પશ્ચિમ અને વાયવ્યની રહેવાની અને પવનની ઝડપ ૧૩ થી ૧૭ કીમી/કલાક રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન ૩૩-૩૪ ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન ૨૩-૨૩ ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૨૮-૦૯-૧૬	૨૯-૦૯-૧૬	૩૦-૦૯-૧૬	૦૧-૧૦-૧૬	૦૨-૧૦-૧૬
વરસાદ(મી.મી)	0	0	0	0	10
મહત્તમ તાપમાન (સે.ગ્રે)	33	34	34	34	34
લઘુત્તમ તાપમાન (સે.ગ્રે)	23	23	23	23	23
વાદળની સ્થિતિ(ઓકટા)	2	1	1	3	3
મહત્તમ ભેજ (%) સવાર	88	88	89	87	82
લઘુત્તમ ભેજ (%) બપોરબાદ	56	53	53	52	52
પવનની ગતિ (કિમી/કલાક)	16	15	14	13	17
પવનની દિશા (ડીગ્રી)	288	277	283	291	295

સુરેન્દ્રનગર: જીલ્લામાં આગામી તા. ૨૮-૦૯-૨૦૧૬ થી તા. ૦૨-૧૦-૨૦૧૬ ભેજવાળું અને અંશતઃ વાદળછાયું રહેશે. વરસાદની શક્યતા નથી. પવનની દિશા પશ્ચિમની રહેવાની અને પવનની ઝડપ **૧૪ થી ૧૯ કીમી/કલાક** રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન **૩૩-૩૩** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન **૨૨-૨૩** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

તારીખ	દિવસ-૧	દિવસ-૨	દિવસ-૩	દિવસ-૪	દિવસ-૫
પેરામીટર	૨૮-૦૯-૧૬	૨૯-૦૯-૧૬	૩૦-૦૯-૧૬	૦૧-૧૦-૧૬	૦૨-૧૦-૧૬
વરસાદ(મી.મી)	૦	૦	૦	૦	૦
મહત્તમ તાપમાન (સે.ગ્રે)	૩૩	૩૩	૩૩	૩૩	૩૩
લઘુત્તમ તાપમાન (સે.ગ્રે)	૨૩	૨૨	૨૨	૨૨	૨૨
વાદળની સ્થિતિ(ઓકટા)	૧	૦	૦	૨	૨
મહત્તમ ભેજ (%) સવાર	૮૨	૮૪	૮૪	૮૦	૭૧
લઘુત્તમ ભેજ (%) બપોરબાદ	૪૧	૩૭	૩૯	૩૮	૩૮
પવનની ગતિ (કિમી/કલાક)	૧૯	૧૭	૧૬	૧૪	૧૮
પવનની દિશા (ડીગ્રી)	૨૭૫	૨૭૧	૨૭૨	૨૮૦	૨૮૩

૦૨ ઓક્ટોબર સુધી હવામાન ભેજવાળું અને વાદળછાયું રહેશે અને પવનની દિશા પશ્ચિમની રહેવાની અને પવનની ઝડપ **૫ થી ૧૫ કીમી/કલાક** રહેવાની શક્યતા છે. આ સમયગાળામાં મહત્તમ તાપમાન દિવસ દરમિયાન **૩૨-૩૪** ડીગ્રી સેલ્શીયસ જેટલું તેમજ લઘુત્તમ તાપમાન રાત્રી દરમિયાન **૨૪-૨૫** ડીગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે.

- ઓછો વરસાદ હોય ત્યાં જમીનમાં ભેજ જળવાય રહે તે માટે આંતર ખેડ અને નિંદામણ કરવું. શક્ય હોય તો પાકના અવશેષોનું આવરણ અથવા પલ્લાસ્ટીકનું આવરણ કરવું. તેમજ તીવ્ર ભેજની અછત હોય ત્યાં ૨ ટકા પોટેશીયમ નાઈટ્રેટના દ્રાવણનો છંટકાવ કરવો.

મુખ્ય પાક	પાક અવસ્થા	રોગ/જીવાત/જાત	કૃષિ સલાહ (૧૦ લીટર પાણીમાં મિશ્ર કરી ભલામણ કરેલ દવા છાંટવી)
મગફળી	સુયા/પોપટા	સફેદ મુંડા  ચીતરી અને ગેરુ  	ફોરેટ ૧૦ ટકા દાણાદાર દવા ૨૫ કી.ગ્રા. પ્રતિ હેક્ટરે જમીનમાં પુરતો ભેજ હોય ત્યારે નાખવી અથવા ક્લોરોપાયરીફોસ દવા ૪ લીટર પ્રતિ હેક્ટરે પિયત પાણી સાથે અથવા ક્લોરોપાયરીફોસ દવા ૨૫ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી જમીનમાં થડનીફરતે રેડવું. પાનના ટપકાના રોગના આગોતરા નિયંત્રણ માટે હેક્ઝાકોનાઝોલ ૧૫ મિલી અથવા તેબુકોનાઝોલ ૧૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.

કપાસ	ચાપવા અને ફૂલ	લીલા તડતડીયા/થ્રીપ્સ 	નિયંત્રણ માટે એસીફેટ ૧૫ ગ્રામ અને એ બામેક્ટીન ૧૦ મી.લી. ૧૦ લીટર પાણીમાં મિશ્ર કરી છાંટવી.
એરંડા	વૃદ્ધિ અને ફુલ	ઘોડિયા ઈ યળ અને લશ્કરી ઈયળ→ 	તેના નિયંત્રણ માટે કક્વીનાલફોસ ૨૦ મીલી ૧૦ લી.પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
ભીંડા	વૃદ્ધિ/ફુલ/ શીંગ	શીંગ કોરી ખાનાર ઈયળ કથીરી	➤ તેના ફળ અને ડોકા કોરી ખાનાર ઈયળનો ઉપદ્રવ વધુ હોય તો સ્પીનોસાડ ૩ મિલી /૧૦ લીટર પાણીમાં ઓગાળી છંટકાવ કરવો. ➤ પ્રોપરગાઈટ ૧૦ મિલી અથવા ડાયકોફોલ ૨૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
તુરીયા, દુધી, ગલકા અને કારેલા	વૃદ્ધિ/ફુલ	પાન કોરીયું , થ્રીપ્સ, સફેદ માખી તલછારો	➤ લીંબોળીનો અર્ક ૫૦૦ મિલી અથવા લીંબોળીનું તેલ ૫૦ મીલીનો છંટકાવ ૧૫ દિવસના અંતરે ચાર વખત કરવો. વધુ ઉપદ્રવ હોય તો એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ / ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. ➤ મેન્કોઝેબ ૨૭ ગ્રામ અથવા ક્લોરોથેલોનીલ ૨૭ ગ્રામ અથવા ફોઝેટાઈલ ૨૦ ગ્રામના ૧૫ દિવસના અંતરે ત્રણ છંટકાવ કરવા.
ગુવાર	વૃદ્ધિ/ફુલ/ શીંગ	થ્રીપ્સ,સફેદ માખી	➤ તેના નિયંત્રણ માટે એસીટામાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અને લીંબોળીનું તેલ ૫૦ મીલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મીલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.

રીંગણી અને ટમેટી	ફુલ/ફળ	થ્રીપ્સ, સફેદ માખી, ફળ અને ડોકા કોરી ખાનાર ઈયળ	➤ લીંબોળીનું તેલ ૫૦ મિલી અથવા લીંબોળી ના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. જો વધુ ઉપદ્રવ હોય તો એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો. રીંગણી અને ટમેટીમાં ફળ અને ડોકા કોરી ખાનાર ઈયળનો ઉપદ્રવ વધુ હોય તો સ્પીનોસાડ ૩ મિલી /૧૦ લીટર પાણીમાં ઓગાળી છંટકાવ કરવો.
મરચી	ફુલ/ફળ	થ્રીપ્સ 	➤ પ્રોફેનોફોસ ૨૦ મિલી અને લીંબોળી નું તેલ ૫૦ મિલી અથવા લીંબોળી ના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી ૧૦ લી. પાણીમાં મિશ્ર કરી છંટકાવ કરવો. 
કાકડી	વૃદ્ધિ/ફુલ/ફળ	થ્રીપ્સ, સફેદ માખી	➤ તેના નિયંત્રણ માટે એસીટેમાપ્રીડ ૨૦ ટકા ૫ ગ્રામ અને લીંબોળીનું તેલ ૫૦ મિલી અથવા લીંબોળીના મીંજનું ૫ ટકા અર્કનું દ્રાવણ ૫૦૦ મિલી/૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો.
લીંબુ	ફળવિકાસ	બદામી ટપકા પાનકોરીયું	કોપરઓક્સીક્લોરાઈડ ૪૦ ગ્રામ અને ૧.૫ ગ્રામ સ્ટ્રેપ્ટોસાયકલીનના બે છંટકાવ ૧૫ દિવસના ગાળે કરવા ડીડીવીપી ૧૦ મી.લી. ૧૦ લી. પાણીમાં મિશ્ર કરી છાંટવી.
5X1qDZ 3F પશુ/મર ઘા	પ્રજીવથી થતા રોગોની ચકાસણી કરાવવી. કરમિયાની દવા આપવી તથા ઉછરતા પશુઓમાં બૃસેલ્લા કાફ્વુંડનું રસીકરણ કરાવવું. ઈતરડીના નિયંત્રણ માટે ડેલ્ટામેથ્રીન અથવા એમીટ્રાઝ ૨ મી.લી. ૧ લીટર પાણીમાં નાખીને છાંટવી. ઉછરતા પશુઓમાં કરમીયાની દવા આપવી અને કાફ્વુંડ અને બૃસેલ્લાનું રસીકરણ કરાવવું. પશુને ૬-૮ કી.ગ્રા. સુકો અને ૧૫-૨૦ કી.ગ્રા. લીલ્લો ચારો આપવો. રહેથાની જગ્યાની સફાઈ રાખવી અને માખી- મચ્છરથી બચવા ફિનાઈલ છાંટવું. 		

પ્રતિ,
તંત્રીશ્રી/ડાયરેક્ટરશ્રી,
રાજકોટ,

ઉપરોક્ત માહિતી આપના દૈનિક સમાચાર પત્રોમાં, રેડીયોમાં ગામના ચો રાના કાર્યક્રમમાં તથા ટીવીના ગ્રામ્ય કાર્યક્રમમાં વિના મૂલ્યે પ્રસિધ્ધ કરાવવા નમ્ર વિનંતી છે. સહકારની અપેક્ષા સહ આભાર.

ટેક્નીકલ ઓફિસર
સુકી ખેતી સંશોધન કેન્દ્ર
જુનાગઢ કૃષિ યુનિવર્સિટી
તરઘડીયા



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-52
for the Rajkot district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
(Period 28 Sept. to 02 Oct. - 2016)

Significant past weather for the preceding week of Rajkot

Rainfall (mm)	0.0
Maximum temperature (°C)	32.0-32.8
Minimum temperature (°C)	21.5-23.8
Relative Humidity (%) I	84-89
Relative Humidity (%) II	58-64
Wind speed (Kmph)	6.8-8.1

Weather forecast until 08-30 hrs of 02/10/2016

(1) RAJKOT DISTRICT:

	28/09/16	29/09/16	30/09/16	01/10/16	02/10/16
Rainfall (mm)	0	0	0	0	10
Max.Temp.(°C)	33	33	33	33	33
Min.Temp.(°C)	23	22	22	22	22
Total CC (octa)	2	1	1	2	2
RH-I (%)	90	91	90	89	83
RH-II (%)	53	50	51	49	49
Wind Speed (kmph)	18	18	15	12	20
Wind Direction(deg.)	282	276	276	286	287

Agro meteorological Advisories:

General Advice	• Rain fall: Humid and mainly cloudy weather forecasted for next five days. Light rainfall may occur on 2.10.2016 • Temperature: Maximum and minimum temperature likely to be in range of 33-33°C and 22-23°C, respectively. • Relative Humidity: The RH is likely to be in the range of 49-90 % • Wind Direction and Speed: Direction and speed of wind is likely to be W and 12-20 km/h, respectively. • Remove weeds and Wherever, possible mulching should be practiced by using dry grass, crop residues, etc. to conserve soil moisture, apply irrigation to the crops as per availability and crop stage. In areas of severe dry spell take a spray of 2% potassium nitrate (2kg in 100 lit. of water) on the standing crop.
-----------------------	--

Crops	Stage	Insect/Pest/Variety	Advisory
Kharif groundnut	Pod Development	White grub  Tikka & rust 	Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha. For prevention of leaf spot disease spray Hexaconazole 10 ml OR Tebuconazole 7 ml in 10 liters of water under clear sky.
Cotton	Squaring /flowering/ball Formation-development	Jassid/Thrips 	Spray acephate 75 % SP@ 15 gm and abamectin 10 ml / 10 lit of water.
Castor	Flowering/capsule formation/development	Semi loopier & prodenia→ 	Heavy infestation of semilooper observed in castor, Spray Quinalphos 20 ml in 10 liters of water.
Lady finger	Flowering/fruiting	Shoot and fruit borer Mites	Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) Spray Propargite 10 ml or Dicofol 20 ml in 10 litres of water. For control of mites.
Ridge gourd/ Bottle gourd/ Sponge gourd/ Bitter gourd	Flowering/Fruiting	White fly Downy mildew & Alternaria blight 	Spray neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosetyl 20 g/10 lit of water at 15 days interval. 
Cluster bean	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval
Brinjal and tomato	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at

			15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips → 	In present weather conditions severe attack of thrips has been reported. Spray spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and 20 ml profenofos in 10 litres of water.
Cucumbers	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval
Livestock & poultry 	Do Protozoan disease evaluation and its treatments. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid fly's and mosquitoes.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-52 for the Jamnagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 28 Sept. to 02 Oct. - 2016)

Weather forecast until 08-30 hrs of 02/10/2016

(2) JAMNAGAR DISTRICT:

	28/09/16	29/09/16	30/09/16	01/10/16	02/10/16
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	32	32	32	32	32
Min.Temp.(°C)	24	24	24	24	24
Total CC (octa)	2	1	0	2	1
RH-I (%)	89	89	88	88	83
RH-II (%)	57	55	56	55	55
Wind Speed (kmph)	18	19	15	11	20
Wind Direction(deg.)	276	270	268	280	285

Agro meteorological Advisories:-

General Advice	• Rain fall : Humid and clear to mainly cloudy weather and no rainfall forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 32-32°C and 24-24°C, respectively. • Relative Humidity: The RH is likely to be in the range of 55-89 % • Wind Direction and Speed: Direction and speed of wind is likely to be W and 11-20 km/h, respectively. • Remove weeds and Wherever, possible mulching should be practiced by using dry grass, crop residues, etc. to conserve soil moisture, apply irrigation to the crops as per availability and crop stage. In areas of severe dry spell take a spray of 2% potassium nitrate (2kg in 100 lit. of water) on the standing crop.		
Crops	Stage	Insect/Pest/Variety	Advisory
Kharif groundnut	Pod formation/Development	White grub  Tikka & rust 	Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha. For prevention of leaf spot disease spray Hexaconazole 10 ml OR Tebuconazole 7 ml in 10 liters of water under clear sky.

Cotton	Squaring /flowering/ball Formation-developmnt	Jassid/Thrips 	Spray acephate 75 % SP@ 15 gm and abamectin 10 ml / 10 lit of water.
Castor	Flowering/capsule formation	Semi looper & prodenia→ 	Heavy infestation of semilooper observed in castor, Spray Quinalphos 20 ml in 10 liters of water.
Lady finger	Flowerin/fruiting	Shoot and fruit borer Mites	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) Spray Propargite 10 ml or Dicofol 20 ml in 10 litres of water. For control of mites.
Ridge gourd/ Bottle gourd/ Sponge gourd/ Bitter gourd	Flowering/Fruiting	White fly Downy mildew & Alternaria blight 	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Cluster bean	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval
Brinjal and tomato	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips → 	In present weather conditions severe attack of thrips has been reported. Spray spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and 20 ml profenofos in 10 litres of water.
Cucumbers	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval

**Livestock
& poultry**



Do Protozoan disease evaluation and its treatments. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid fly's and mosquitoes.

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell :94277 28523



Agro met Advisory Service bulletin No.-52 for the Amreli district
 (Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)
 (Period 28 Sept. to 02 Oct. - 2016)

Weather forecast until 08-30 hrs of 02/10/2016

(3) AMRELI DISTRICT:

	28/09/16	29/09/16	30/09/16	01/10/16	02/10/16
Rainfall (mm)	0	0	0	0	10
Max.Temp.(°C)	33	34	34	34	34
Min.Temp.(°C)	23	23	23	23	23
Total CC (octa)	2	1	1	3	3
RH-I (%)	88	88	89	87	82
RH-II (%)	56	53	53	52	52
Wind Speed (kmph)	16	15	14	13	17
Wind Direction(deg.)	288	277	283	291	295

Agro meteorological Advisories:

General Advice	• Rain fall: Humid and mainly to partly cloudy weather forecasted for next five days and light rainfall on 02 October forecasted. • Temperature: Maximum and minimum temperature likely to be in range of 33-34°C and 23-23°C, respectively. • Relative Humidity: The RH is likely to be in the range of 52-89 % • Wind Direction and Speed: Direction and speed of wind is likely to be W, NW and 13-16 km/h, respectively. • Remove weeds and Wherever, possible mulching should be practiced by using dry grass, crop residues, etc. to conserve soil moisture, apply irrigation to the crops as per availability and crop stage. In areas of severe dry spell take a spray of 2% potassium nitrate (2kg in 100 lit. of water) on the standing crop.			
Crops	Stage	Insect/Pest/Variety	Advisory	
Kharif groundnut	Pod formation/Development	White grub  Tikka & rust 	Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha. For prevention of leaf spot disease spray Hexaconazole 10 ml OR Tebuconazole 7 ml in 10 liters of water under clear sky.	

Cotton	Squaring /flowering/ball Formation-developmmt	Jassid/Thrips 	Spray acephate 75 % SP@ 15 gm and abamectin 10 ml / 10 lit of water.
Castor	Flowering/capsule formation	Semi looper & prodenia→ 	Heavy infestation of semilooper observed in castor, Spray Quinalphos 20 ml in 10 liters of water.
Lady finger	Flowerin/ fruiting	Shoot and fruit borer Mites	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) Spray Propargite 10 ml or Dicofol 20 ml in 10 litres of water. For control of mites.
Ridge gourd/ Bottle gourd/ Sponge gourd/ Bitter gourd	Flowering/Fruiting	White fly Downy mildew & Alternaria blight 	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Cluster bean	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval
Brinjal and tomato	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips → 	In present weather conditions severe attack of thrips has been reported. Spray spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and 20 ml profenofos in 10 litres of water.
Cucumbers	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at

			15 days interval
Livestock & poultry 	Do Protozoan disease evaluation and its treatments. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid fly's and mosquitoes.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.



Gramin Krushi Mausam Seva
Main Dry Farming Research Station
Junagadh Agricultural University
Targhadia (Rajkot) - 360 003
Phone: (0281)2784260(O), 2331686(R)
Fax : (0281)2784722, Cell:94277 28523



Agro met Advisory Service bulletin No.-52 for the Surendranagar district

(Issued jointly by DFRS, J.A.U., Targhadia (Rajkot) and India Meteorological Department)

(Period 28 Sept. to 02 Oct. - 2016)


Weather forecast until 08-30 hrs of 02/10/2016

(4) SURENDRANAGAR DISTRICT:

	28/09/16	29/09/16	30/09/16	01/10/16	02/10/16
Rainfall (mm)	0	0	0	0	0
Max.Temp.(°C)	33	33	33	33	33
Min.Temp.(°C)	23	22	22	22	22
Total CC (octa)	1	0	0	2	2
RH-I (%)	82	84	84	80	71
RH-II (%)	41	37	39	38	38
Wind Speed (kmph)	19	17	16	14	18
Wind Direction(deg.)	275	271	272	280	283

Agro meteorological advisories:

General Advice	• Rain fall: Humid and clear to mainly cloudy weather and no rainfall forecasted for next five days. • Temperature: Maximum and minimum temperature likely to be in range of 33-33°C and 22-22°C, respectively. • Relative Humidity: The RH is likely to be in the range of 37-84 % • Wind Direction and Speed: Direction and speed of wind is likely to be W and 14-19 km/h, respectively. • Remove weeds and Wherever, possible mulching should be practiced by using dry grass, crop residues, etc. to conserve soil moisture, apply irrigation to the crops as per availability and crop stage. In areas of severe dry spell take a spray of 2% potassium nitrate (2kg in 100 lit. of water) on the standing crop.		
Crops	Stage	Insect/Pest/Variety	Advisory
Kharif groundnut	Pod formation/Development	White grub  Tikka & rust 	Do soil application of Phorate 10 G 25 kg/ha or Drenching of Chlorpyrifos or Quinalphos 4 liters/ha. For prevention of leaf spot disease spray Hexaconazole 10 ml OR Tebuconazole 7 ml in 10 liters of water under clear sky.

Cotton	Squaring /flowering/ball Formation-developmmt	Jassid/Thrips 	Spray acephate 75 % SP@ 15 gm and abamectin 10 ml / 10 lit of water.
Castor	Flowering/capsule formation	Semi looper & prodenia→ 	Heavy infestation of semilooper observed in castor, Spray Quinalphos 20 ml in 10 liters of water.
Lady finger	Flowerin/ fruiting	Shoot and fruit borer Mites	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval(4 spray) Spray Propargite 10 ml or Dicofol 20 ml in 10 litres of water. For control of mites.
Ridge gourd/ Bottle gourd/ Sponge gourd/ Bitter gourd	Flowering/Fruiting	White fly Downy mildew & Alternaria blight 	Spary neem seed extract 500 ml or neem seed oil 50 ml in 10 litre of water at 15 days interval (4 sprays) or acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. Alternate three spray of Mancozeb 27 g or chlorothalonil 27 g or fosatyl 20 g/10 lit of water at 15 days interval. 
Cluster bean	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval
Brinjal and tomato	Growing/fruiting	Sucking pest, Fruit and shoot borer	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at 15 days interval. To control shoot and fruit borer, infested fruits and shoots should be collected and buried inside the soil. Installation of pheromone trap @ 2-3 traps per acre is advised for monitoring of borers. If pest population is high, spraying of Spinosad 48 EC @ 1 ml /4 litres of water is advised when sky remains clear.
Chili	Flowering & fruiting	Thrips → 	In present weather conditions severe attack of thrips has been reported. Spray spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and 20 ml profenofos in 10 litres of water.
Cucumbers	Growing/fruiting	Thrips, white fly	To control them spray neem seed extract 500 ml or neem seed oil 50 or neem oil 40 ml and acetamiprid 20 % SP@ 5.0 gm / 10 lit of water at

			15 days interval.
Livestock & poultry 	Do Protozoan disease evaluation and its treatments. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. Spray phenyl in the animal shed to avoid fly's and mosquitoes.		

Technical Officer, GKMS Unit, DFRS, JAU, Targhadia.